

6-2017

Order, Distance, Closure and Convergence: Reconciling Competing Fundamental Topological Concepts

Walter Tholen

York University, tholen@mathstat.yorku.ca

Follow this and additional works at: http://ecommons.udayton.edu/topology_conf



Part of the [Geometry and Topology Commons](#), and the [Special Functions Commons](#)

eCommons Citation

Tholen, Walter, "Order, Distance, Closure and Convergence: Reconciling Competing Fundamental Topological Concepts" (2017).
Summer Conference on Topology and Its Applications. 49.
http://ecommons.udayton.edu/topology_conf/49

This Plenary Lecture is brought to you for free and open access by the Department of Mathematics at eCommons. It has been accepted for inclusion in Summer Conference on Topology and Its Applications by an authorized administrator of eCommons. For more information, please contact frice1@udayton.edu, mschlangen1@udayton.edu.

Order, Distance, Closure and Convergence: Reconciling Competing Fundamental Topological Concepts

Walter Tholen

York University, Toronto, Canada

32nd Summer Conference on Topology and Its Applications,
University of Dayton, June 27–30, 2017

Acknowledgements

Thanks to

Dirk Hofmann (University of Aveiro, Portugal)

for his pointers to Hausdorff's book; see:

D. Hofmann, C. Reis: Convergence and quantale-enriched categories,
arXiv(math):1705.08671

Hongliang Lai (Sichuan University, China)

for his contributions to our joint papers pertaining to this talk; see:

H. Lai, W. Tholen: Quantale-valued topological spaces via closure and convergence, *Topology and Its Applications* (to appear)
arXiv(math):1604.08813

H. Lai, W. Tholen: A note on the topologicity of quantale-valued topological spaces, *Logical Methods in Computer Science* (to appear)
arXiv(cs):1612.09504

Convergence and Metric: Maurice Fréchet 1906

“Sur quelques points du calcul fonctionnel”

Rendiconti del Circolo Matematico di Palermo 22, 1–74

Axioms on the convergence of sequences (p 6):

- $(A) \rightarrow A$
- $(A_n) \rightarrow A \implies (A_{n_i}) \rightarrow A$

Axioms on the distance between points (p 18):

- $(A, B) = (B, A) \geq 0$
- $(A, B) = 0 \iff A = B$
- For some function f with $\lim_{\varepsilon > 0} f(\varepsilon) = 0$:
 $(A, B) \leq \varepsilon, (B, C) \leq \varepsilon \implies (A, C) \leq f(\varepsilon)$

Convergence and Metric: Maurice Fréchet 1906

“Sur quelques points du calcul fonctionnel”

Rendiconti del Circolo Matematico di Palermo 22, 1–74

Axioms on the convergence of sequences (p 6):

- $(A) \rightarrow A$
- $(A_n) \rightarrow A \implies (A_{n_i}) \rightarrow A$

Axioms on the distance between points (p 18):

- $(A, B) = (B, A) \geq 0$
- $(A, B) = 0 \iff A = B$
- For some function f with $\lim_{\varepsilon > 0} f(\varepsilon) = 0$:
 $(A, B) \leq \varepsilon, (B, C) \leq \varepsilon \implies (A, C) \leq f(\varepsilon)$

Convergence and Metric: Maurice Fréchet 1906

“Sur quelques points du calcul fonctionnel”

Rendiconti del Circolo Matematico di Palermo 22, 1–74

Axioms on the convergence of sequences (p 6):

- $(A) \rightarrow A$
- $(A_n) \rightarrow A \implies (A_{n_i}) \rightarrow A$

Axioms on the distance between points (p 18):

- $(A, B) = (B, A) \geq 0$
- $(A, B) = 0 \iff A = B$
- For some function f with $\lim_{\varepsilon > 0} f(\varepsilon) = 0$:
 $(A, B) \leq \varepsilon, (B, C) \leq \varepsilon \implies (A, C) \leq f(\varepsilon)$

Metric-Topology-Convergence: Felix Hausdorff 1914

“Grundzüge der Mengenlehre” (Veit & Comp., Leipzig), Ch. 7, p 211:

“Welchen der drei [...] Grundbegriffe Entfernung, Limes, Umgebung man zur Basis der Betrachtung wählen will, ist bis zu einem gewissen Grade Geschmacksache. [...] Danach scheint die Entfernungstheorie die speziellste, die Limestheorie die allgemeinste zu sein; auf der andern Seite bringt der Limesbegriff sofort eine Beziehung zum Abzählbaren (zu Elementfolgen) in die Theorie hinein, worauf die Umgebungstheorie verzichtet.”

“Which of the three [...] fundamental notions, distance, limit, neighbourhood, one wants to choose as the basis of consideration is, to a certain degree, a matter of taste. [...] Accordingly, the distance theory seems to be the most special, the limit theory the most general; on the other side, the notion of limit brings immediately a connection to countability (to sequences of elements), which the neighbourhood theory foregoes.”

“Grundzüge der Mengenlehre” (Veit & Comp., Leipzig), Ch. 7, p 211:

“Welchen der drei [...] Grundbegriffe Entfernung, Limes, Umgebung man zur Basis der Betrachtung wählen will, ist bis zu einem gewissen Grade Geschmacksache. [...] Danach scheint die Entfernungstheorie die speziellste, die Limestheorie die allgemeinste zu sein; auf der andern Seite bringt der Limesbegriff sofort eine Beziehung zum Abzählbaren (zu Elementfolgen) in die Theorie hinein, worauf die Umgebungstheorie verzichtet.”

“Which of the three [...] fundamental notions, distance, limit, neighbourhood, one wants to choose as the basis of consideration is, to a certain degree, a matter of taste. [...] Accordingly, the distance theory seems to be the most special, the limit theory the most general; on the other side, the notion of limit brings immediately a connection to countability (to sequences of elements), which the neighbourhood theory foregoes.”

Haudorff's functional approach to ...

- Order:

$$M \times M \longrightarrow \{<, >, =\}, \quad M \times M \longrightarrow \{<, >, =, ||\}$$

- Metric:

$$M \times M \longrightarrow \mathbb{R}$$

- Convergence:

$$M^{\mathbb{N}} \dashrightarrow M \quad (\iff M^{\mathbb{N}} \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$

- Topology (via neighbourhood systems):

$$M \longrightarrow 2^{2^M} \quad (\iff 2^M \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$
$$x \mapsto \mathcal{U}(x)$$

Haudorff's functional approach to ...

- Order:

$$M \times M \longrightarrow \{<, >, =\}, \quad M \times M \longrightarrow \{<, >, =, ||\}$$

- Metric:

$$M \times M \longrightarrow \mathbb{R}$$

- Convergence:

$$M^{\mathbb{N}} \dashrightarrow M \quad (\iff M^{\mathbb{N}} \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$

- Topology (via neighbourhood systems):

$$M \longrightarrow 2^{2^M} \quad (\iff 2^M \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$
$$x \mapsto \mathcal{U}(x)$$

Haudorff's functional approach to ...

- Order:

$$M \times M \longrightarrow \{<, >, =\}, \quad M \times M \longrightarrow \{<, >, =, ||\}$$

- Metric:

$$M \times M \longrightarrow \mathbb{R}$$

- Convergence:

$$M^{\mathbb{N}} \dashrightarrow M \quad (\iff M^{\mathbb{N}} \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$

- Topology (via neighbourhood systems):

$$M \longrightarrow 2^{2^M} \quad (\iff 2^M \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$
$$x \mapsto \mathcal{U}(x)$$

Haudorff's functional approach to ...

- Order:

$$M \times M \longrightarrow \{<, >, =\}, \quad M \times M \longrightarrow \{<, >, =, ||\}$$

- Metric:

$$M \times M \longrightarrow \mathbb{R}$$

- Convergence:

$$M^{\mathbb{N}} \dashrightarrow M \quad (\iff M^{\mathbb{N}} \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$

- Topology (via neighbourhood systems):

$$M \longrightarrow 2^{2^M} \quad (\iff 2^M \times M \longrightarrow 2 = \{\text{true}, \text{false}\})$$
$$x \mapsto \mathcal{U}(x)$$

“Nun steht einer Verallgemeinerung dieser Vorstellung nichts im Wege, und wir können uns denken, daß eine beliebige Funktion der Paare einer Menge definiert, d. h. jedem Paar (a, b) von Elementen einer Menge M ein bestimmtes Element $n = f(a, b)$ einer zweiten Menge N zugeordnet sei. In noch weiterer Verallgemeinerung können wir eine Funktion der Elementtripel, Elementfolgen, Elementkomplexe, Teilmengen u. dgl. von M in Betracht ziehen.”

“Now there is no obstacle to generalizing this point of view, and we could think that there be defined an arbitrary function of pairs of a set, i.e., that, to every pair (a, b) of elements in a set M , there be assigned a certain element $n = f(a, b)$ in a second set N . In even further generalization we can consider a function of triples of elements, sequences of elements, complexes [= families] of elements, subsets of M , and the like.”

... but he also cautions:

“Eine ganz allgemein gehaltene Theorie dieser Art würde natürlich erhebliche Komplikationen bedingen und wenig positive Ausbeute liefern.”

“If kept very general, a theory of this type would of course cause considerable complications and provide little positive outcome.”

OUR GENERAL STRATEGY (to keep useless generality in check):
Put structure on Hausdorff's set N which can capture the syntax used in key axioms of the theory!

BUT WE MUST DECIDE EARLY ON:
Keep N fixed, or let N “float” (together with M)?
Guided by our principal examples, we'll keep it fixed, but we'll briefly return to this point at the end of this talk!

... but he also cautions:

“Eine ganz allgemein gehaltene Theorie dieser Art würde natürlich erhebliche Komplikationen bedingen und wenig positive Ausbeute liefern.”

“If kept very general, a theory of this type would of course cause considerable complications and provide little positive outcome.”

OUR GENERAL STRATEGY (to keep useless generality in check):
Put structure on Hausdorff’s set N which can capture the syntax used in key axioms of the theory!

BUT WE MUST DECIDE EARLY ON:

Keep N fixed, or let N “float” (together with M)?

Guided by our principal examples, we’ll keep it fixed, but we’ll briefly return to this point at the end of this talk!

... but he also cautions:

“Eine ganz allgemein gehaltene Theorie dieser Art würde natürlich erhebliche Komplikationen bedingen und wenig positive Ausbeute liefern.”

“If kept very general, a theory of this type would of course cause considerable complications and provide little positive outcome.”

OUR GENERAL STRATEGY (to keep useless generality in check):
Put structure on Hausdorff’s set N which can capture the syntax used in key axioms of the theory!

BUT WE MUST DECIDE EARLY ON:
Keep N fixed, or let N “float” (together with M)?
Guided by our principal examples, we’ll keep it fixed, but we’ll briefly return to this point at the end of this talk!

Seeing a metric as “hom”: F. William Lawvere 1973

Folklore since Eilenberg–Mac Lane in the 1940s:

Preordered sets	\Leftrightarrow	“thin” (small) categories
$\text{true} \Rightarrow x \leq x$		$1 \longrightarrow \text{hom}(x, x)$
$x \leq y \ \& \ y \leq z \Rightarrow x \leq z$		$\text{hom}(x, y) \times \text{hom}(y, z) \longrightarrow \text{hom}(x, z)$

“Metric spaces, generalized logic, and closed categories”,
Rendiconti del Seminario Matematico e Fisico di Milano 43, 135–166

$0 \geq d(x, x)$
 $d(x, y) + d(y, z) \geq d(x, z)$
“homs” are just numbers!

“Syntax providers”:

$\mathbf{2} = ((\{\text{true}, \text{false}\}, \Rightarrow), \&, \text{true})$ $\mathbf{Set} = ((\{\text{all sets}\}, \longrightarrow), \times, 1)$
 $\underline{[0, \infty]} = (([0, \infty], \geq), +, 0)$

Seeing a metric as “hom”: F. William Lawvere 1973

Folklore since Eilenberg–Mac Lane in the 1940s:

Preordered sets	\Leftrightarrow	“thin” (small) categories
$\text{true} \Rightarrow x \leq x$		$1 \longrightarrow \text{hom}(x, x)$
$x \leq y \ \& \ y \leq z \Rightarrow x \leq z$		$\text{hom}(x, y) \times \text{hom}(y, z) \longrightarrow \text{hom}(x, z)$

“Metric spaces, generalized logic, and closed categories”,

Rendiconti del Seminario Matematico e Fisico di Milano 43, 135–166

$$0 \geq d(x, x)$$

$$d(x, y) + d(y, z) \geq d(x, z)$$

“homs” are just numbers!

“Syntax providers”:

$$\mathbf{2} = ((\{\text{true}, \text{false}\}, \Rightarrow), \&, \text{true})$$

$$\mathbf{[0, \infty]} = (([0, \infty], \geq), +, 0)$$

$$\mathbf{Set} = ((\{\text{all sets}\}, \longrightarrow), \times, 1)$$

Seeing a metric as “hom”: F. William Lawvere 1973

Folklore since Eilenberg–Mac Lane in the 1940s:

Preordered sets	\Leftrightarrow	“thin” (small) categories
$\text{true} \Rightarrow x \leq x$		$1 \longrightarrow \text{hom}(x, x)$
$x \leq y \ \& \ y \leq z \Rightarrow x \leq z$		$\text{hom}(x, y) \times \text{hom}(y, z) \longrightarrow \text{hom}(x, z)$

“Metric spaces, generalized logic, and closed categories”,
Rendiconti del Seminario Matematico e Fisico di Milano 43, 135–166

$0 \geq d(x, x)$
 $d(x, y) + d(y, z) \geq d(x, z)$
“homs” are just numbers!

“Syntax providers”:

$\mathbf{2} = ((\{\text{true}, \text{false}\}, \Rightarrow), \&, \text{true})$ $\mathbf{Set} = ((\{\text{all sets}\}, \longrightarrow), \times, 1)$
 $\mathbf{[0, \infty]} = (([0, \infty], \geq), +, 0)$

What's special about the extra structure?

Cartesian closedness, exponentiability, monoidal closedness:

$$\frac{Z \times X \longrightarrow Y}{Z \longrightarrow Y^X}$$

Set

$$(\sum Z_i) \times X \cong \sum(Z_i \times X)$$

$$\frac{Z \times X \longrightarrow Y}{Z \longrightarrow C(X, Y)}$$

???**Top**???

$$(\sum Z_i) \times X \cong \sum(Z_i \times X)$$

$$\frac{C \otimes A \longrightarrow B}{C \longrightarrow \text{hom}(A, B)}$$

AbGrp

$$(\bigoplus C_i) \otimes A \cong \bigoplus(C_i \otimes A)$$

$$\frac{z + x \geq y}{z \geq \max\{y-x, 0\}}$$

$\overleftarrow{[0, \infty]}$

$$(\inf z_i) + x = \inf(z_i + x)$$

V unital (but not necessarily commutative) *quantale*
= complete lattice with a monoid structure $V = (V, \otimes, k)$ s.th.

$$u \otimes \bigvee_{i \in I} v_i = \bigvee_{i \in I} u \otimes v_i, \quad \left(\bigvee_{i \in I} v_i \right) \otimes u = \bigvee_{i \in I} (v_i \otimes u)$$

- $V = 2$ with $u \otimes v = u \wedge v$, $k = \text{true}$
- $V = \overleftarrow{[0, \infty]}$ with $u \otimes v = u + v$, $k = 0$
- V any frame with $u \otimes v = u \wedge v$, $k = \top$
- $V = 2^M$ (M monoid) with $A \otimes B = \{\alpha\beta \mid \alpha \in A, \beta \in B\}$, $k = \{\eta_M\}$

“Internal hom” operations:

$$v \leq w \swarrow u \iff v \otimes u \leq w \iff u \leq v \searrow w$$

V unital (but not necessarily commutative) *quantale*
= complete lattice with a monoid structure $V = (V, \otimes, k)$ s.th.

$$u \otimes \bigvee_{i \in I} v_i = \bigvee_{i \in I} u \otimes v_i, \quad \left(\bigvee_{i \in I} v_i \right) \otimes u = \bigvee_{i \in I} (v_i \otimes u)$$

- $V = 2$ with $u \otimes v = u \wedge v$, $k = \text{true}$
- $V = \overleftarrow{[0, \infty]}$ with $u \otimes v = u + v$, $k = 0$
- V any frame with $u \otimes v = u \wedge v$, $k = \top$
- $V = 2^M$ (M monoid) with $A \otimes B = \{\alpha\beta \mid \alpha \in A, \beta \in B\}$, $k = \{\eta_M\}$

“Internal hom” operations:

$$v \leq w \swarrow u \iff v \otimes u \leq w \iff u \leq v \searrow w$$

V unital (but not necessarily commutative) *quantale*
 = complete lattice with a monoid structure $V = (V, \otimes, k)$ s.th.

$$u \otimes \bigvee_{i \in I} v_i = \bigvee_{i \in I} u \otimes v_i, \quad \left(\bigvee_{i \in I} v_i \right) \otimes u = \bigvee_{i \in I} (v_i \otimes u)$$

- $V = 2$ with $u \otimes v = u \wedge v$, $k = \text{true}$
- $V = \overleftarrow{[0, \infty]}$ with $u \otimes v = u + v$, $k = 0$
- V any frame with $u \otimes v = u \wedge v$, $k = \top$
- $V = 2^M$ (M monoid) with $A \otimes B = \{\alpha\beta \mid \alpha \in A, \beta \in B\}$, $k = \{\eta_M\}$

“Internal hom” operations:

$$v \leq w \swarrow u \iff v \otimes u \leq w \iff u \leq v \searrow w$$

Quantale-enriched categories

(V, \otimes, k)

V-Cat:

V-category :

$(X, a : X \times X \rightarrow V) :$ $k \leq a(x, x)$
 $a(y, z) \otimes a(x, y) \leq a(x, z)$

V-functor :

$(X, a) \xrightarrow{f} (Y, b) :$ $a(x, y) \leq b(fx, fy)$

$V = 2$

V-Cat = Ord = {(pre)ordered sets}

$V = \overleftarrow{[0, \infty]}$

V-Cat = Met = {(gen'd) metric spaces}

$V = 2^M$
(for a monoid M)

V-Cat = M-Ord = {" M -ordered sets"} :

$$X \leq_{\eta_M} X$$

$$X \leq_{\alpha} Y, Y \leq_{\beta} Z \Rightarrow X \leq_{\beta\alpha} Z$$

Quantale-enriched categories

(V, \otimes, k)

V-Cat:

V-category:

$(X, a : X \times X \rightarrow V) :$ $k \leq a(x, x)$
 $a(y, z) \otimes a(x, y) \leq a(x, z)$

V-functor:

$(X, a) \xrightarrow{f} (Y, b) :$ $a(x, y) \leq b(fx, fy)$

$V = 2$

V-Cat = Ord

= {(pre)ordered sets}

$V = \overleftarrow{[0, \infty]}$

V-Cat = Met

= {(gen'd) metric spaces}

$V = 2^M$
(for a monoid M)

V-Cat = M-Ord

= {"M-ordered sets"} :

$$X \leq_{\eta_M} X$$

$$X \leq_{\alpha} Y, Y \leq_{\beta} Z \Rightarrow X \leq_{\beta\alpha} Z$$

Quantale-enriched categories

(V, \otimes, k)

V-Cat:

V-category :

$(X, a : X \times X \rightarrow V) :$ $k \leq a(x, x)$
 $a(y, z) \otimes a(x, y) \leq a(x, z)$

V-functor :

$(X, a) \xrightarrow{f} (Y, b) :$ $a(x, y) \leq b(fx, fy)$

$V = 2$

V-Cat = Ord

= {(pre)ordered sets}

$V = \overleftarrow{[0, \infty]}$

V-Cat = Met

= {(gen'd) metric spaces}

$V = 2^M$
(for a monoid M)

V-Cat = M-Ord

= {"M-ordered sets"} :

$$X \leq_{\eta_M} X$$

$$X \leq_{\alpha} Y, Y \leq_{\beta} Z \Rightarrow X \leq_{\beta\alpha} Z$$

Quantale-enriched categories

(V, \otimes, k)

V-Cat:

V-category :

$(X, a : X \times X \rightarrow V) :$ $k \leq a(x, x)$
 $a(y, z) \otimes a(x, y) \leq a(x, z)$

V-functor :

$(X, a) \xrightarrow{f} (Y, b) :$ $a(x, y) \leq b(fx, fy)$

$V = 2$

V-Cat = Ord

= {(pre)ordered sets}

$V = \overleftarrow{[0, \infty]}$

V-Cat = Met

= {(gen'd) metric spaces}

$V = 2^M$
(for a monoid M)

V-Cat = M -Ord

= {" M -ordered sets"} :

$$X \leq_{\eta_M} X$$

$$X \leq_{\alpha} Y, Y \leq_{\beta} Z \Rightarrow X \leq_{\beta\alpha} Z$$

Quantale-enriched categories

(V, \otimes, k)

V-Cat:

V-category :

$(X, a : X \times X \rightarrow V) :$ $k \leq a(x, x)$
 $a(y, z) \otimes a(x, y) \leq a(x, z)$

V-functor :

$(X, a) \xrightarrow{f} (Y, b) :$ $a(x, y) \leq b(fx, fy)$

$V = 2$

V-Cat = Ord

= {(pre)ordered sets}

$V = \overleftarrow{[0, \infty]}$

V-Cat = Met

= {(gen'd) metric spaces}

$V = 2^M$
(for a monoid M)

V-Cat = M-Ord

= {"M-ordered sets"} :

$x \leq_{\eta_M} x$

$x \leq_{\alpha} y, y \leq_{\beta} z \Rightarrow x \leq_{\beta\alpha} z$

One more quantale: {distance distribution functions}

$$\overleftarrow{[0, \infty]} \cong \overrightarrow{[0, 1]} = (([0, 1], \leq), \cdot, 1)$$

$$\Delta \ni \varphi : [0, \infty] \longrightarrow [0, 1] \quad \varphi(\beta) = \sup_{\alpha < \beta} \varphi(\alpha)$$

$$(\varphi \otimes \psi)(\gamma) = \sup_{\alpha + \beta = \gamma} \varphi(\alpha) \psi(\beta) \quad \kappa(\alpha) = \begin{cases} 0 & \text{if } \alpha = 0, \\ 1 & \text{if } \alpha > 0. \end{cases}$$

$$\sigma : \overleftarrow{[0, \infty]} \leftrightarrow \Delta \leftrightarrow \overrightarrow{[0, 1]} : \tau$$

$$\varphi = \sup_{\gamma} \sigma(\gamma) \otimes \tau(\varphi(\gamma)) : \quad \Delta \text{ as a coproduct in } \mathbf{CQuant} !$$

$$\mathbf{Met} \leftrightarrow \mathbf{ProbMet} \leftrightarrow \mathbf{ProbOrd} (\cong \mathbf{Met})$$

[Menger 1942], [Schweizer-Sklar 1983]

One more quantale: {distance distribution functions}

$$\overleftarrow{[0, \infty]} \cong \overrightarrow{[0, 1]} = (([0, 1], \leq), \cdot, 1)$$

$$\Delta \ni \varphi : [0, \infty] \longrightarrow [0, 1] \quad \varphi(\beta) = \sup_{\alpha < \beta} \varphi(\alpha)$$

$$(\varphi \otimes \psi)(\gamma) = \sup_{\alpha + \beta = \gamma} \varphi(\alpha) \psi(\beta) \quad \kappa(\alpha) = \begin{cases} 0 & \text{if } \alpha = 0, \\ 1 & \text{if } \alpha > 0. \end{cases}$$

$$\sigma : \overleftarrow{[0, \infty]} \leftrightarrow \Delta \leftrightarrow \overrightarrow{[0, 1]} : \tau$$

$$\varphi = \sup_{\gamma} \sigma(\gamma) \otimes \tau(\varphi(\gamma)) : \quad \Delta \text{ as a coproduct in } \mathbf{CQuant} !$$

$$\mathbf{Met} \leftrightarrow \mathbf{ProbMet} \leftrightarrow \mathbf{ProbOrd} (\cong \mathbf{Met})$$

[Menger 1942], [Schweizer-Sklar 1983]

One more quantale: {distance distribution functions}

$$\overleftarrow{[0, \infty]} \cong \overrightarrow{[0, 1]} = (([0, 1], \leq), \cdot, 1)$$

$$\Delta \ni \varphi : [0, \infty] \longrightarrow [0, 1] \quad \varphi(\beta) = \sup_{\alpha < \beta} \varphi(\alpha)$$

$$(\varphi \otimes \psi)(\gamma) = \sup_{\alpha + \beta = \gamma} \varphi(\alpha) \psi(\beta) \quad \kappa(\alpha) = \begin{cases} 0 & \text{if } \alpha = 0, \\ 1 & \text{if } \alpha > 0. \end{cases}$$

$$\sigma : \overleftarrow{[0, \infty]} \leftrightarrow \Delta \leftrightarrow \overrightarrow{[0, 1]} : \tau$$

$$\varphi = \sup_{\gamma} \sigma(\gamma) \otimes \tau(\varphi(\gamma)) : \quad \Delta \text{ as a coproduct in } \mathbf{CQuant} !$$

$$\mathbf{Met} \leftrightarrow \mathbf{ProbMet} \leftrightarrow \mathbf{ProbOrd} (\cong \mathbf{Met})$$

[Menger 1942], [Schweizer-Sklar 1983]

One more quantale: {distance distribution functions}

$$\overleftarrow{[0, \infty]} \cong \overrightarrow{[0, 1]} = (([0, 1], \leq), \cdot, 1)$$

$$\Delta \ni \varphi : [0, \infty] \longrightarrow [0, 1] \quad \varphi(\beta) = \sup_{\alpha < \beta} \varphi(\alpha)$$

$$(\varphi \otimes \psi)(\gamma) = \sup_{\alpha + \beta = \gamma} \varphi(\alpha) \psi(\beta) \quad \kappa(\alpha) = \begin{cases} 0 & \text{if } \alpha = 0, \\ 1 & \text{if } \alpha > 0. \end{cases}$$

$$\sigma : \overleftarrow{[0, \infty]} \leftrightarrow \Delta \leftrightarrow \overrightarrow{[0, 1]} : \tau$$

$$\varphi = \sup_{\gamma} \sigma(\gamma) \otimes \tau(\varphi(\gamma)) : \quad \Delta \text{ as a coproduct in } \mathbf{CQuant} !$$

$$\mathbf{Met} \leftrightarrow \mathbf{ProbMet} \leftrightarrow \mathbf{ProbOrd} (\cong \mathbf{Met})$$

[Menger 1942], [Schweizer-Sklar 1983]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Now: $a : PX \times X \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

Need $\hat{P}a : PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

Choose: $\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Rewrite: $c : PX \rightarrow 2^X$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\bigcup \mathcal{A})$

Simplify:

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

Equiv'ly:

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -Cat = Cls

closure spaces & continuous maps

[Seal 2005]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Now: $a : PX \times PX \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

Need $\hat{P}a$: $PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

Choose: $\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Rewrite: $c : PX \rightarrow 2^X$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\bigcup \mathcal{A})$

Simplify:

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

Equiv'ly:

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -Cat = Cls

closure spaces & continuous maps

[Seal 2005]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

$f : (X, a) \rightarrow (Y, b)$

Now: $a : PX \times X \rightarrow 2$

Need $\hat{P}a : PPX \times PX \rightarrow 2$

Choose: $\mathbb{P} = (P, \cup, \{-\})$

Rewrite: $c : PX \rightarrow 2^X$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

(R) true $\Rightarrow a(\{x\}, x)$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\cup \mathcal{A}, z)$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\cup \mathcal{A})$

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -Cat = Cls

closure spaces & continuous maps

[Seal 2005]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Now: $a : PX \times X \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

Need $\hat{P}a$: $PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

Choose: $\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Rewrite: $c : PX \rightarrow 2^X$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\bigcup \mathcal{A})$

Simplify:

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

Equiv'ly:

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -Cat = Cls

closure spaces & continuous maps

[Seal 2005]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Now: $a : PX \times X \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

Need $\hat{P}a : PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

Choose: $\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Rewrite: $c : PX \rightarrow 2^X$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\bigcup \mathcal{A})$

Simplify:

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

Equiv'ly:

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -Cat = Cls

closure spaces & continuous maps

[Seal 2005]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Now: $a : PX \times X \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

Need $\hat{P}a : PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

Choose: $\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Rewrite: $c : PX \rightarrow 2^X$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\bigcup \mathcal{A})$

Simplify:

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

Equiv'ly:

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -Cat = Cls

closure spaces & continuous maps

[Seal 2005]

Hausdorff's vision, I: Replace points by subsets

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Now: $a : PX \times X \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

Need $\hat{P}a : PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

Choose: $\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Rewrite: $c : PX \rightarrow 2^X$

(R) $\{x\} \subseteq c\{x\}$

(T) $B \subseteq \bigcup_{A \in \mathcal{A}} cA \ \& \ z \in cB \Rightarrow z \in c(\bigcup \mathcal{A})$

Simplify:

(R*) $A \subseteq cA$

(T*) $B \subseteq cA \ \& \ z \in cB \Rightarrow z \in cA$

Equiv'ly:

$B \subseteq cA \Rightarrow cB \subseteq cA$

c extensive and idempotent!

Also: $f : X \rightarrow Y$

(C*) $f(c_X A) \subseteq c_Y(fA)$

Hence: $(\mathbb{P}, 2)$ -**Cat** = **Cls**
[Seal 2005]

closure spaces & continuous maps

Top

$$c : PX \longrightarrow 2^X$$

$$(R) A \subseteq cA$$

$$(T) B \subseteq cA \Rightarrow cB \subseteq cA$$

c fin'ly additive:

$$(A) c(A \cup B) = cA \cup cB$$

$$c(\emptyset) = \emptyset$$

$$(C) f(c_X A) \subseteq c_Y(fA)$$

[Seal 2009]

V-Cls

$$c : PX \longrightarrow V^X$$

$$(R) \forall x \in A : k \leq (cA)(x)$$

= (\mathbb{P}, V) -Cat

$$(T) \left(\bigwedge_{y \in B} (cA)(y) \right) \otimes (cB)(x) \leq (cA)(x)$$

[Lai-T 2016]

V-Top

c fin'ly additive:

$$(A) c(A \cup B)(x) = (cA)(x) \vee (cB)(x)$$

$$c(\emptyset)(x) = \perp$$

$$(C) (c_X A)(x) \leq c_Y(fA)(fx)$$

(Not to be confused with other V-concepts in topology!)

Cls \leftrightarrow Top, V-Cls \leftrightarrow V-Top

Top

$$c : PX \longrightarrow 2^X$$

$$(R) A \subseteq cA$$

$$(T) B \subseteq cA \Rightarrow cB \subseteq cA$$

c fin'ly additive:

$$(A) c(A \cup B) = cA \cup cB$$

$$c(\emptyset) = \emptyset$$

$$(C) f(c_X A) \subseteq c_Y(fA)$$

[Seal 2009]

V-Cls

$$c : PX \longrightarrow V^X$$

$$(R) \forall x \in A : k \leq (cA)(x)$$

= (\mathbb{P}, V) -Cat

$$(T) \left(\bigwedge_{y \in B} (cA)(y) \right) \otimes (cB)(x) \leq (cA)(x)$$

[Lai-T 2016]

V-Top

c fin'ly additive:

$$(A) c(A \cup B)(x) = (cA)(x) \vee (cB)(x)$$

$$c(\emptyset)(x) = \perp$$

$$(C) (c_X A)(x) \leq c_Y(fA)(fx)$$

(Not to be confused with other V-concepts in topology!)

$$V = \overleftarrow{[0, \infty]}, \Delta, \dots$$

$\overleftarrow{[0, \infty]}$ -Cls:

$$\delta : X \times \mathcal{P}X \longrightarrow \overleftarrow{[0, \infty]} \quad (\text{R}) \quad \forall x \in A : 0 \geq \delta(x, A)$$

$$\delta(x, A) = (cA)(x) \quad (\text{T}) \quad (\sup_{y \in B} \delta(y, A)) + \delta(x, B) \geq \delta(x, A)$$

$\overleftarrow{[0, \infty]}$ -Top = App [Lowen 1997] (but his condition (T) is different!):

δ finitely additive

$$(\text{A}) \quad \delta(x, A \cup B) = \min\{\delta(x, A), \delta(x, B)\}$$

$$\delta(x, \emptyset) = \infty$$

$$f : X \longrightarrow Y \quad (\text{C}) \quad \delta_X(x, A) \geq \delta_Y(fx, fA)$$

Δ -Top = ProbApp [Jäger 2015]

$\Delta_{\&}$ -Top, with $\&$ denoting any left-continuous t-norm on $[0,1]$ –
all “generated” by only three such norms!

$$V = \overleftarrow{[0, \infty]}, \Delta, \dots$$

$\overleftarrow{[0, \infty]}$ -Cls:

$$\delta : X \times \mathcal{P}X \longrightarrow \overleftarrow{[0, \infty]} \quad (\text{R}) \quad \forall x \in A : 0 \geq \delta(x, A)$$

$$\delta(x, A) = (cA)(x) \quad (\text{T}) \quad (\sup_{y \in B} \delta(y, A)) + \delta(x, B) \geq \delta(x, A)$$

$\overleftarrow{[0, \infty]}$ -Top = App [Lowen 1997] (but his condition (T) is different!):

δ finitely additive

$$(\text{A}) \quad \delta(x, A \cup B) = \min\{\delta(x, A), \delta(x, B)\}$$

$$\delta(x, \emptyset) = \infty$$

$$f : X \longrightarrow Y$$

$$(\text{C}) \quad \delta_X(x, A) \geq \delta_Y(fx, fA)$$

Δ -Top = ProbApp [Jäger 2015]

$\Delta_{\&}$ -Top, with $\&$ denoting any left-continuous t-norm on $[0,1]$ –
all “generated” by only three such norms!

$$V = \overleftarrow{[0, \infty]}, \Delta, \dots$$

$\overleftarrow{[0, \infty]}$ -Cls:

$$\delta : X \times \mathcal{P}X \longrightarrow \overleftarrow{[0, \infty]} \quad (\text{R}) \quad \forall x \in A : 0 \geq \delta(x, A)$$

$$\delta(x, A) = (cA)(x) \quad (\text{T}) \quad (\sup_{y \in B} \delta(y, A)) + \delta(x, B) \geq \delta(x, A)$$

$\overleftarrow{[0, \infty]}$ -Top = App [Lowen 1997] (but his condition (T) is different!):

δ finitely additive $(\text{A}) \quad \delta(x, A \cup B) = \min\{\delta(x, A), \delta(x, B)\}$

$$\delta(x, \emptyset) = \infty$$

$$f : X \longrightarrow Y \quad (\text{C}) \quad \delta_X(x, A) \geq \delta_Y(fx, fA)$$

Δ -Top = ProbApp [Jäger 2015]

$\Delta_{\&}$ -Top, with $\&$ denoting any left-continuous t-norm on $[0,1]$ –
all “generated” by only three such norms!

$$V = \overleftarrow{[0, \infty]}, \Delta, \dots$$

$\overleftarrow{[0, \infty]}$ -Cls:

$$\delta : X \times \mathcal{P}X \longrightarrow \overleftarrow{[0, \infty]} \quad (\text{R}) \quad \forall x \in A : 0 \geq \delta(x, A)$$

$$\delta(x, A) = (cA)(x) \quad (\text{T}) \quad (\sup_{y \in B} \delta(y, A)) + \delta(x, B) \geq \delta(x, A)$$

$\overleftarrow{[0, \infty]}$ -Top = App [Lowen 1997] (but his condition (T) is different!):

δ finitely additive $(\text{A}) \quad \delta(x, A \cup B) = \min\{\delta(x, A), \delta(x, B)\}$

$$\delta(x, \emptyset) = \infty$$

$$f : X \longrightarrow Y$$

$$(\text{C}) \quad \delta_X(x, A) \geq \delta_Y(fx, fA)$$

Δ -Top = ProbApp [Jäger 2015]

$\Delta_{\&}$ -Top, with $\&$ denoting any left-continuous t-norm on $[0, 1]$ –
all “generated” by only three such norms!

Properties: Is V-Top topological over Set?

V-CIs certainly is:

Initial structure c on X w.r.t. $(f_i : X \rightarrow (Y_i, c_i), i \in I)$ (I any size)
(= largest c making all f_i continuous):

$$(cA)(x) = \bigwedge_{i \in I} c_i(f_i A)(f_i x)$$

But: $(\forall i \in I : c_i \text{ finitely additive}) \not\Rightarrow c \text{ finitely additive}$

Fix:

$$(c^+ A)(x) = \bigwedge_{A=M_1 \cup \dots \cup M_n} (cM_1)(x) \vee \dots \vee (cM_n)(x)$$

Did (T) survive the passage $c \mapsto c^+$?

Properties: Is V-Top topological over Set?

V-CIs certainly is:

Initial structure c on X w.r.t. $(f_i : X \rightarrow (Y_i, c_i), i \in I)$ (I any size)
(= largest c making all f_i continuous):

$$(cA)(x) = \bigwedge_{i \in I} c_i(f_i A)(f_i x)$$

But: $(\forall i \in I : c_i \text{ finitely additive}) \not\Rightarrow c \text{ finitely additive}$

Fix:

$$(c^+ A)(x) = \bigwedge_{A=M_1 \cup \dots \cup M_n} (cM_1)(x) \vee \dots \vee (cM_n)(x)$$

Did (T) survive the passage $c \mapsto c^+$?

“More distributivity” needed for V !

[Dikranjan-T 1995]: $\overline{c^+}$ is finitely additive; hence:

$$\overline{c^+} \leq \bar{c} \leq c \implies \overline{c^+} \leq c^+$$

But what should \bar{c} be here?

$$(\bar{c}A)(x) := \bigvee_{v \in V} v \otimes c(c^v A)(x), \quad \text{with } c^v A := \{z \in X \mid v \leq (cA)(z)\}$$

If $v = p$ **coprime** ($p > \perp$, $p \leq u \vee v \Rightarrow p \leq u$ or $p \leq v$):

$$c^p(A \cup B) = c^p A \cup c^p B, \quad c^p \emptyset = \emptyset$$

Need:

$$V = \{\bigvee p_i \mid p_i \text{ coprime}\},$$

i.e. $V \cong \{\text{closed sets}\}$ for some topological space X : V **spatial coframe**

“More distributivity” needed for V !

[Dikranjan-T 1995]: $\overline{c^+}$ is finitely additive; hence:

$$\overline{c^+} \leq \overline{c} \leq c \implies \overline{c^+} \leq c^+$$

But what should \overline{c} be here?

$$(\overline{c}A)(x) := \bigvee_{v \in V} v \otimes c(c^v A)(x), \quad \text{with } c^v A := \{z \in X \mid v \leq (cA)(z)\}$$

If $v = p$ **coprime** ($p > \perp$, $p \leq u \vee v \Rightarrow p \leq u$ or $p \leq v$):

$$c^p(A \cup B) = c^p A \cup c^p B, \quad c^p \emptyset = \emptyset$$

Need:

$$V = \{\bigvee p_i \mid p_i \text{ coprime}\},$$

i.e. $V \cong \{\text{closed sets}\}$ for some topological space X : V **spatial coframe**

“More distributivity” needed for V !

[Dikranjan-T 1995]: $\overline{c^+}$ is finitely additive; hence:

$$\overline{c^+} \leq \overline{c} \leq c \implies \overline{c^+} \leq c^+$$

But what should \overline{c} be here?

$$(\overline{c}A)(x) := \bigvee_{v \in V} v \otimes c(c^v A)(x), \quad \text{with } c^v A := \{z \in X \mid v \leq (cA)(z)\}$$

If $v = p$ **coprime** ($p > \perp$, $p \leq u \vee v \Rightarrow p \leq u$ or $p \leq v$):

$$c^p(A \cup B) = c^p A \cup c^p B, \quad c^p \emptyset = \emptyset$$

Need:

$$V = \{\bigvee p_i \mid p_i \text{ coprime}\},$$

i.e. $V \cong \{\text{closed sets}\}$ for some topological space X : V **spatial coframe**

“More distributivity” needed for V !

[Dikranjan-T 1995]: $\overline{c^+}$ is finitely additive; hence:

$$\overline{c^+} \leq \overline{c} \leq c \implies \overline{c^+} \leq c^+$$

But what should \overline{c} be here?

$$(\overline{c}A)(x) := \bigvee_{v \in V} v \otimes c(c^v A)(x), \quad \text{with } c^v A := \{z \in X \mid v \leq (cA)(z)\}$$

If $v = p$ **coprime** ($p > \perp$, $p \leq u \vee v \Rightarrow p \leq u$ or $p \leq v$):

$$c^p(A \cup B) = c^p A \cup c^p B, \quad c^p \emptyset = \emptyset$$

Need:

$$V = \{\bigvee p_i \mid p_i \text{ coprime}\},$$

i.e. $V \cong \{\text{closed sets}\}$ for some topological space X : V **spatial coframe**

Topologicity of **V-Top** over **Set**

Conclusion:

V spatial coframe \implies V-Top (bi)coreflective in V-CIs

Status of the hypothesis?

Without invoking Choice:

V continuous, spatial coframe \implies V constructively completely distributive

i.e. $\bigvee : \text{Down}V \longrightarrow V$ preserves all infima

“ \Leftarrow ” needs Choice, and then $(\text{ccd} \iff \text{cd})$

[Gierz *et al.* 2003], [Wood 2004]

Topologicity of **V-Top** over **Set**

Conclusion:

V spatial coframe \implies **V-Top** (bi)coreflective in **V-CIs**

Status of the hypothesis?

Without invoking Choice:

V continuous, spatial coframe \implies V constructively completely distributive

i.e. $\bigvee : \text{Down}V \longrightarrow V$ preserves all infima

“ \Leftarrow ” needs Choice, and then $(\text{ccd} \iff \text{cd})$

[Gierz *et al.* 2003], [Wood 2004]

Topologicity of **V-Top** over **Set**

Conclusion:

V spatial coframe \implies **V-Top** (bi)coreflective in **V-CIs**

Status of the hypothesis?

Without invoking Choice:

V continuous, spatial coframe \implies V constructively completely distributive

i.e. $\bigvee : \text{Down}V \longrightarrow V$ preserves all infima

“ \Leftarrow ” needs Choice, and then $(\text{ccd} \iff \text{cd})$

[Gierz *et al.* 2003], [Wood 2004]

Hausdorff's vision, II: Replace points by (ultra)filters

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Also: $a : PX \times PX \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

$\hat{P}a : PPX \times PPX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

$\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Now: $a : UX \times UX \rightarrow 2$

(R) true $\Rightarrow a(\dot{x}, x)$

$\bar{U}a : UUX \times UUX \rightarrow 2$

(T) $\bar{U}a(\mathfrak{X}, \eta) \ \& \ a(\eta, z) \Rightarrow a(\Sigma \mathfrak{X}, z)$

$\mathbb{U} = (U, \Sigma, (-))$

$\bar{U}a(\mathfrak{X}, \eta) : \iff \forall A \in \mathfrak{X}, B \in \eta$
 $\exists \mathfrak{x} \in \mathcal{A}, y \in B : a(\mathfrak{x}, y)$

Again:

(R) $\dot{x} \rightarrow x$

(T) $\mathfrak{x} \rightarrow \eta \ \& \ \eta \rightarrow z \Rightarrow \Sigma \mathfrak{x} \rightarrow z$

$f : (X, \rightarrow) \rightarrow (Y, \rightarrow)$

(C) $\mathfrak{x} \rightarrow y \Rightarrow f[\mathfrak{x}] \rightarrow fy$

[Barr 1970]: $(\mathbb{U}, 2)$ -Cat \cong Top

[Seal 2005]: Works also for filters!

Hausdorff's vision, II: Replace points by (ultra)filters

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Also: $a : PX \times PX \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

$\hat{P}a : PPX \times PX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

$\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Now: $a : UX \times UX \rightarrow 2$

(R) true $\Rightarrow a(\dot{x}, x)$

$\bar{U}a : UUX \times UX \rightarrow 2$

(T) $\bar{U}a(\mathfrak{X}, \eta) \ \& \ a(\eta, z) \Rightarrow a(\Sigma \mathfrak{X}, z)$

$\mathbb{U} = (U, \Sigma, (-))$

$\bar{U}a(\mathfrak{X}, \eta) : \iff \forall A \in \mathfrak{X}, B \in \eta$
 $\exists \mathfrak{r} \in \mathcal{A}, y \in B : a(\mathfrak{r}, y)$

Again:

(R) $\dot{x} \rightarrow x$

(T) $\mathfrak{X} \rightarrow \eta \ \& \ \eta \rightarrow z \Rightarrow \Sigma \mathfrak{X} \rightarrow z$

$f : (X, \rightarrow) \rightarrow (Y, \rightarrow)$

(C) $\mathfrak{r} \rightarrow y \Rightarrow f[\mathfrak{r}] \rightarrow fy$

[Barr 1970]: $(\mathbb{U}, 2)$ -Cat \cong Top

[Seal 2005]: Works also for filters!

Hausdorff's vision, II: Replace points by (ultra)filters

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Also: $a : PX \times PX \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

$\hat{P}a : PPX \times PPX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

$\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Now: $a : UX \times UX \rightarrow 2$

(R) true $\Rightarrow a(\dot{x}, x)$

$\bar{U}a : UUX \times UUX \rightarrow 2$

(T) $\bar{U}a(\mathfrak{X}, \eta) \ \& \ a(\eta, z) \Rightarrow a(\Sigma \mathfrak{X}, z)$

$\mathbb{U} = (U, \Sigma, (-))$

$\bar{U}a(\mathfrak{X}, \eta) : \iff \forall A \in \mathfrak{X}, B \in \eta$
 $\exists \mathfrak{x} \in \mathcal{A}, y \in B : a(\mathfrak{x}, y)$

Again:

(R) $\dot{x} \rightarrow x$

(T) $\mathfrak{x} \rightarrow \eta \ \& \ \eta \rightarrow z \Rightarrow \Sigma \mathfrak{x} \rightarrow z$

$f : (X, \rightarrow) \rightarrow (Y, \rightarrow)$

(C) $\mathfrak{x} \rightarrow y \Rightarrow f[\mathfrak{x}] \rightarrow fy$

[Barr 1970]: $(\mathbb{U}, 2)\text{-Cat} \cong \mathbf{Top}$

[Seal 2005]: Works also for filters!

Hausdorff's vision, II: Replace points by (ultra)filters

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Also: $a : PX \times PX \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

$\hat{P}a : PPX \times PPX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

$\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Now: $a : UX \times UX \rightarrow 2$

(R) true $\Rightarrow a(\dot{x}, x)$

$\bar{U}a : UUX \times UUX \rightarrow 2$

(T) $\bar{U}a(\mathfrak{X}, \eta) \ \& \ a(\eta, z) \Rightarrow a(\Sigma \mathfrak{X}, z)$

$\mathbb{U} = (U, \Sigma, (-))$

$\bar{U}a(\mathfrak{X}, \eta) : \iff \forall A \in \mathfrak{X}, B \in \eta$

$\exists x \in A, y \in B : a(x, y)$

Again:

(R) $\dot{x} \rightarrow x$

(T) $\mathfrak{X} \rightarrow \eta \ \& \ \eta \rightarrow z \Rightarrow \Sigma \mathfrak{X} \rightarrow z$

$f : (X, \rightarrow) \rightarrow (Y, \rightarrow)$

(C) $x \rightarrow y \Rightarrow f[x] \rightarrow fy$

[Barr 1970]: $(\mathbb{U}, 2)\text{-Cat} \cong \mathbf{Top}$

[Seal 2005]: Works also for filters!

Hausdorff's vision, II: Replace points by (ultra)filters

Recall: $a : X \times X \rightarrow 2$

(R) true $\Rightarrow a(x, x)$

(T) $a(x, y) \ \& \ a(y, z) \Rightarrow a(x, z)$

$f : (X, a) \rightarrow (Y, b)$

(C) $a(x, y) \Rightarrow b(fx, fy)$

Also: $a : PX \times PX \rightarrow 2$

(R) true $\Rightarrow a(\{x\}, x)$

$\hat{P}a : PPX \times PPX \rightarrow 2$

(T) $\hat{P}a(\mathcal{A}, B) \ \& \ a(B, z) \Rightarrow a(\bigcup \mathcal{A}, z)$

$\mathbb{P} = (P, \bigcup, \{-\})$

$\hat{P}a(\mathcal{A}, B) : \iff \forall y \in B \exists A \in \mathcal{A} : a(A, y)$

Now: $a : UX \times UX \rightarrow 2$

(R) true $\Rightarrow a(\dot{x}, x)$

$\bar{U}a : UUX \times UUX \rightarrow 2$

(T) $\bar{U}a(\mathfrak{X}, \eta) \ \& \ a(\eta, z) \Rightarrow a(\Sigma \mathfrak{X}, z)$

$\mathbb{U} = (U, \Sigma, (-))$

$\bar{U}a(\mathfrak{X}, \eta) : \iff \forall A \in \mathfrak{X}, B \in \eta$
 $\exists x \in A, y \in B : a(x, y)$

Again:

(R) $\dot{x} \rightarrow x$

(T) $\mathfrak{X} \rightarrow \eta \ \& \ \eta \rightarrow z \Rightarrow \Sigma \mathfrak{X} \rightarrow z$

$f : (X, \rightarrow) \rightarrow (Y, \rightarrow)$

(C) $x \rightarrow y \Rightarrow f[x] \rightarrow fy$

[Barr 1970]: $(\mathbb{U}, 2)\text{-Cat} \cong \mathbf{Top}$

[Seal 2005]: Works also for filters!

$(\mathbb{U}, \mathbb{V})\text{-Cat} \cong \mathbb{V}\text{-Top} ?$

\mathbb{V} completely distributive

For $a : \mathbb{U}X \times X \rightarrow \mathbb{V}$, define $\bar{\mathbb{U}}a : \mathbb{U}\mathbb{U}X \times \mathbb{U}X \rightarrow \mathbb{V}$:

$$\bar{\mathbb{U}}a(\mathfrak{x}, \mathfrak{y}) := \bigwedge_{A \in \mathfrak{x}, B \in \mathfrak{y}} \bigvee_{x \in A, y \in B} a(x, y)$$

\mathbb{P} and \mathbb{U} interact via the \mathbb{V} -relation $\varepsilon_X : \mathbb{P}X \times \mathbb{U}X \rightarrow \mathbb{V}$:

$$\varepsilon_X(A, \mathfrak{x}) = \begin{cases} k & \text{if } A \in \mathfrak{x} \\ \perp & \text{else} \end{cases} \quad A_\varepsilon : (\mathbb{U}, \mathbb{V})\text{-Cat} \rightarrow (\mathbb{P}, \mathbb{V})\text{-Cat}$$
$$(X, a) \mapsto (X, c_a = a \circ \varepsilon_X)$$

$$(c_a A)(y) = \bigvee_{x \ni A} a(x, y)$$

A_ε has a right adjoint $R : (X, c) \mapsto (X, a_c)$, $a_c(x, y) = \bigwedge_{A \in \mathfrak{x}} (cA)(y)$

$(\mathbb{U}, \mathbb{V})\text{-Cat} \cong \mathbb{V}\text{-Top} ?$

\mathbb{V} completely distributive

For $a : \mathbb{U}X \times X \rightarrow \mathbb{V}$, define $\bar{\mathbb{U}}a : \mathbb{U}\mathbb{U}X \times \mathbb{U}X \rightarrow \mathbb{V}$:

$$\bar{\mathbb{U}}a(\mathfrak{x}, \mathfrak{y}) := \bigwedge_{A \in \mathfrak{x}, B \in \mathfrak{y}} \bigvee_{x \in A, y \in B} a(x, y)$$

\mathbb{P} and \mathbb{U} interact via the \mathbb{V} -relation $\varepsilon_X : \mathbb{P}X \times \mathbb{U}X \rightarrow \mathbb{V}$:

$$\varepsilon_X(A, \mathfrak{x}) = \begin{cases} k & \text{if } A \in \mathfrak{x} \\ \perp & \text{else} \end{cases} \quad A_\varepsilon : (\mathbb{U}, \mathbb{V})\text{-Cat} \rightarrow (\mathbb{P}, \mathbb{V})\text{-Cat}$$
$$(X, a) \mapsto (X, c_a = a \circ \varepsilon_X)$$

$$(c_a A)(y) = \bigvee_{x \ni A} a(x, y)$$

A_ε has a right adjoint $R : (X, c) \mapsto (X, a_c)$, $a_c(x, y) = \bigwedge_{A \in \mathfrak{x}} (cA)(y)$

$(\mathbb{U}, \mathbb{V})\text{-Cat} \cong \mathbb{V}\text{-Top} ?$

\mathbb{V} completely distributive

For $a : \mathbb{U}X \times X \rightarrow \mathbb{V}$, define $\bar{\mathbb{U}}a : \mathbb{U}\mathbb{U}X \times \mathbb{U}X \rightarrow \mathbb{V}$:

$$\bar{\mathbb{U}}a(\mathfrak{x}, \mathfrak{y}) := \bigwedge_{A \in \mathfrak{x}, B \in \mathfrak{y}} \bigvee_{x \in A, y \in B} a(x, y)$$

\mathbb{P} and \mathbb{U} interact via the \mathbb{V} -relation $\varepsilon_X : \mathbb{P}X \times \mathbb{U}X \rightarrow \mathbb{V}$:

$$\varepsilon_X(A, \mathfrak{x}) = \begin{cases} k & \text{if } A \in \mathfrak{x} \\ \perp & \text{else} \end{cases} \quad A_\varepsilon : (\mathbb{U}, \mathbb{V})\text{-Cat} \rightarrow (\mathbb{P}, \mathbb{V})\text{-Cat}$$
$$(X, a) \mapsto (X, c_a = a \circ \varepsilon_X)$$

$$(c_a A)(y) = \bigvee_{x \ni A} a(x, y)$$

$$A_\varepsilon \text{ has a right adjoint } R : (X, c) \mapsto (X, a_c), \quad a_c(\mathfrak{x}, y) = \bigwedge_{A \in \mathfrak{x}} (cA)(y)$$

Theorem: Yes !

[Lai-T 2016] V completely distributive. Then:

- $A_\varepsilon : (\mathbb{U}, V)\text{-Cat} \hookrightarrow (\mathbb{P}, V)\text{-Cat} = V\text{-Cls}$ coreflective embedding
- Its image is $V\text{-Top} \cong (\mathbb{U}, V)\text{-Cat}$.

Topologicity of $V\text{-Top}$ revisited:

$(\mathbb{U}, V)\text{-Cat}$ is (almost) trivially topological over \mathbf{Set} !

Corollaries:

[Clementino-Hofmann 2003]

$\mathbf{App} \cong (\mathbb{U}, \overleftarrow{[0, \infty]})\text{-Cat}$

$0 \geq a(\dot{x}, x)$

$(\sup_{A \in \mathcal{X}} \inf_{r \in A} a(r, y)) + a(\eta, z) \geq a(\Sigma \mathcal{X}, z)$

[Jäger 2016],[Lai-T 2016]

$\mathbf{ProbApp} \cong (\mathbb{U}, \Delta)\text{-Cat}$

$(R) \kappa \leq a(\dot{x}, x)$

(T) ...

Theorem: Yes !

[Lai-T 2016] V completely distributive. Then:

- $A_\varepsilon : (\mathbb{U}, V)\text{-Cat} \hookrightarrow (\mathbb{P}, V)\text{-Cat} = V\text{-Cls}$ coreflective embedding
- Its image is $V\text{-Top} \cong (\mathbb{U}, V)\text{-Cat}$.

Topologicity of $V\text{-Top}$ revisited:

$(\mathbb{U}, V)\text{-Cat}$ is (almost) trivially topological over **Set** !

Corollaries:

[Clementino-Hofmann 2003]

$\mathbf{App} \cong (\mathbb{U}, \overleftarrow{[0, \infty]})\text{-Cat}$

$0 \geq a(\dot{x}, x)$

$(\sup_{A \in \mathcal{X}} \inf_{r \in A} a(r, y)) + a(\eta, z) \geq a(\Sigma \mathcal{X}, z)$

[Jäger 2016],[Lai-T 2016]

$\mathbf{ProbApp} \cong (\mathbb{U}, \Delta)\text{-Cat}$

$(R) \kappa \leq a(\dot{x}, x)$

(T) ...

Theorem: Yes!

[Lai-T 2016] V completely distributive. Then:

- $A_\varepsilon : (\mathbb{U}, V)\text{-Cat} \hookrightarrow (\mathbb{P}, V)\text{-Cat} = V\text{-Cls}$ coreflective embedding
- Its image is $V\text{-Top} \cong (\mathbb{U}, V)\text{-Cat}$.

Topologicity of $V\text{-Top}$ revisited:

$(\mathbb{U}, V)\text{-Cat}$ is (almost) trivially topological over **Set**!

Corollaries:

[Clementino-Hofmann 2003]

$\mathbf{App} \cong (\mathbb{U}, \overleftarrow{[0, \infty]})\text{-Cat}$

$0 \geq a(\dot{x}, x)$

$(\sup_{A \in \mathcal{X}} \inf_{r \in A} a(r, y)) + a(\eta, z) \geq a(\Sigma \mathcal{X}, z)$

[Jäger 2016],[Lai-T 2016]

$\mathbf{ProbApp} \cong (\mathbb{U}, \Delta)\text{-Cat}$

$(R) \kappa \leq a(\dot{x}, x)$

(T) ...

Change-of-base functors

\mathbf{V} with $k = \top > \perp$, compl. distributive. Then $\mathbf{2} \leftrightarrow \mathbf{V}$ has both adjoints:

$$\mathbf{2} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{V}$$

Expect transfer of this double adjunction to the categories of models:

$$\mathbf{Ord} = \mathbf{2}\text{-Cat} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{V}\text{-Cat}$$

Moreover,

$$\mathbf{2} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} [0, \infty] \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \Delta$$

$$\mathbf{Ord} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{Met} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{ProbMet}$$

$$\mathbf{Top} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{App} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{ProbApp}$$

Actually: lower row “sits” above upper row, via the functors induced by

$$\begin{array}{c} \mathbf{I} \\ \downarrow (-) \\ \mathbf{U} \end{array}$$

Change-of-base functors

V with $k = \top > \perp$, compl. distributive. Then $2 \leftrightarrow V$ has both adjoints:

$$2 \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} V$$

Expect transfer of this double adjunction to the categories of models:

$$\mathbf{Ord} = \mathbf{2-Cat} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{V-Cat}$$

Moreover,

$$2 \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} [0, \infty] \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \Delta$$

$$\mathbf{Ord} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{Met} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{ProbMet}$$

$$\mathbf{Top} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{App} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{ProbApp}$$

Actually: lower row “sits” above upper row, via the functors induced by

$$\begin{array}{c} \mathbb{I} \\ \downarrow (-) \\ \mathbb{U} \end{array}$$

Change-of-base functors

V with $k = \top > \perp$, compl. distributive. Then $2 \leftrightarrow V$ has both adjoints:

$$2 \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} V$$

Expect transfer of this double adjunction to the categories of models:

$$\mathbf{Ord} = \mathbf{2-Cat} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{V-Cat}$$

Moreover,

$$2 \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} [0, \infty] \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \Delta$$

$$\mathbf{Ord} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{Met} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{ProbMet}$$

$$\mathbf{Top} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{App} \begin{array}{c} \leftarrow \\ \rightleftarrows \\ \rightarrow \end{array} \mathbf{ProbApp}$$

Actually: lower row “sits” above upper row, via the functors induced by

$$\begin{array}{c} \mathbb{I} \\ \downarrow (-) \\ \mathbb{U} \end{array}$$

Change-of-base functors

V with $k = \top > \perp$, compl. distributive. Then $2 \leftrightarrow V$ has both adjoints:

$$2 \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} V$$

Expect transfer of this double adjunction to the categories of models:

$$\mathbf{Ord} = \mathbf{2-Cat} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{V-Cat}$$

Moreover,

$$2 \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} [0, \infty] \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \Delta$$

$$\mathbf{Ord} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{Met} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{ProbMet}$$

$$\mathbf{Top} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{App} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{ProbApp}$$

Actually: lower row “sits” above upper row, via the functors induced by

$$\begin{array}{c} \mathbb{I} \\ \downarrow (-) \\ \mathbb{U} \end{array}$$

Change-of-base functors

V with $k = \top > \perp$, compl. distributive. Then $2 \leftrightarrow V$ has both adjoints:

$$2 \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} V$$

Expect transfer of this double adjunction to the categories of models:

$$\mathbf{Ord} = \mathbf{2-Cat} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{V-Cat}$$

Moreover,

$$2 \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} [0, \infty] \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \Delta$$

$$\mathbf{Ord} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{Met} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{ProbMet}$$

$$\mathbf{Top} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{App} \begin{array}{c} \longleftarrow \\ \longrightarrow \\ \longleftarrow \\ \longrightarrow \end{array} \mathbf{ProbApp}$$

Actually: lower row “sits” above upper row, via the functors induced by

$$\begin{array}{c} \mathbb{I} \\ \downarrow (-) \\ \mathbb{U} \end{array}$$

[Lai-T 2016] V, W completely distributive and integral ($k = \top$),
 $\varphi : V \longrightarrow W$ monotone, $\psi : W \longrightarrow V$ lax homomorphism of quantales.

Equivalent are:

- $\varphi \dashv \psi : W \longrightarrow V$
- $\overline{B_\varphi} \dashv B_\psi : \mathbf{W-Cat} \longrightarrow \mathbf{V-Cat}$, $(X, b) \mapsto (X, \psi b)$
- $\overline{B_\varphi} \dashv B_\psi : (\mathbb{U}, \mathbf{W})\text{-Cat} \longrightarrow (\mathbb{U}, \mathbf{V})\text{-Cat}$

[Here $\overline{B_\varphi}$ assigns to a V -category (X, a) the W -category (X, b) , with the least W -category structure with $b \geq \varphi a$; likewise when (X, a) is an (\mathbb{U}, W) -category.]


(\mathbb{T}, V) -categories: “Topology = Lax Algebra”

$\mathbb{T} = (T, m: TT \rightarrow T, e: Id \rightarrow T)$ **Set**-monad, $\hat{T}: V\text{-Rel} \rightarrow V\text{-Rel}$ lax ext.

$(\mathbb{T}, V)\text{-Cat}$: $(X, a: TX \dashrightarrow X)$: $k \leq a(e_X(x), x)$

$\hat{T}a: TTX \dashrightarrow TX$: $a(\eta, z) \otimes \hat{T}a(x, \eta) \leq a(m_X(x), z)$

$(X, a) \xrightarrow{f} (Y, b)$: $a(x, y) \leq b(Tf(x), f(y))$



Lax Eilenberg-Moore via (Kleisli)^{op}: $e_X^\circ \leq a$
 $a \circ a \leq a$


(\mathbb{T}, V) -categories: “Topology = Lax Algebra”

$\mathbb{T} = (T, m: TT \rightarrow T, e: Id \rightarrow T)$ **Set**-monad, $\hat{T}: V\text{-Rel} \rightarrow V\text{-Rel}$ lax ext.

$(\mathbb{T}, V)\text{-Cat}$: $(X, a: TX \dashrightarrow X)$: $k \leq a(e_X(x), x)$

$\hat{T}a: TTX \dashrightarrow TX$: $a(\eta, z) \otimes \hat{T}a(x, \eta) \leq a(m_X(x), z)$

$(X, a) \xrightarrow{f} (Y, b)$: $a(x, y) \leq b(Tf(x), f(y))$



Lax Eilenberg-Moore via (Kleisli)^{op}: $e_X^\circ \leq a$
 $a \circ a \leq a$




(\mathbb{T}, V) -categories: “Topology = Lax Algebra”

$\mathbb{T} = (T, m: TT \rightarrow T, e: Id \rightarrow T)$ **Set**-monad, $\hat{T}: V\text{-Rel} \rightarrow V\text{-Rel}$ lax ext.

(\mathbb{T}, V) -**Cat**: $(X, a: TX \dashrightarrow X)$: $k \leq a(e_X(x), x)$

$\hat{T}a: TTX \dashrightarrow TX$: $a(\eta, z) \otimes \hat{T}a(x, \eta) \leq a(m_X(x), z)$

$(X, a) \xrightarrow{f} (Y, b)$: $a(x, y) \leq b(Tf(x), f(y))$



Lax Eilenberg-Moore via (Kleisli)^{op}: $e_X^o \leq a$
 $a \circ a \leq a$




(\mathbb{T}, V) -categories: “Topology = Lax Algebra”

$\mathbb{T} = (T, m: TT \rightarrow T, e: Id \rightarrow T)$ **Set-monad**, $\hat{T}: V\text{-Rel} \rightarrow V\text{-Rel}$ lax ext.

$(\mathbb{T}, V)\text{-Cat}$: $(X, a: TX \dashrightarrow X)$: $k \leq a(e_X(x), x)$

$\hat{T}a: TTX \dashrightarrow TX$: $a(\eta, z) \otimes \hat{T}a(x, \eta) \leq a(m_X(x), z)$

$(X, a) \xrightarrow{f} (Y, b)$: $a(x, y) \leq b(Tf(x), f(y))$



Lax Eilenberg-Moore via (Kleisli)^{op}: $e_X^\circ \leq a$
 $a \circ a \leq a$

Hausdorff's vision, III: "What about my nbhd systems?"

$$(X, \mathcal{U}: X \rightarrow \mathbf{FX}) \quad \begin{array}{l} \{-\} \leq \mathcal{U} \\ \mathcal{U} \circ \mathcal{U} \leq \mathcal{U} \end{array} \quad \begin{array}{l} \forall A \in \mathcal{U}(x) : x \in A \\ \forall A \in \mathcal{U}(x) \exists B \in \mathcal{U}(x) \forall y \in B : A \in \mathcal{U}(y) \end{array}$$

[Gähler *et al.* 1992]

[Seal 2005]

Topological spaces are monoids in the (ordered) Kleisli category of the filter monad $\mathbb{F} = (F, \Sigma, \{-\})$ on **Set**:

Top $\cong (\mathbb{F}, 2)$ -**Cat**

More generally:

T-Mon $\cong (\mathbb{T}, 2)$ -**Cat**, for \mathbb{T} "power-enriched"

Hausdorff's vision, III: "What about my nbhd systems?"

$$(X, \mathcal{U}: X \rightarrow \mathbf{FX}) \quad \begin{array}{l} \{-\} \leq \mathcal{U} \\ \mathcal{U} \circ \mathcal{U} \leq \mathcal{U} \end{array} \quad \begin{array}{l} \forall A \in \mathcal{U}(x) : x \in A \\ \forall A \in \mathcal{U}(x) \exists B \in \mathcal{U}(x) \forall y \in B : A \in \mathcal{U}(y) \end{array}$$

[Gähler *et al.* 1992]

[Seal 2005]

Topological spaces are monoids in the (ordered) Kleisli category of the filter monad $\mathbb{F} = (\mathbb{F}, \Sigma, \{-\})$ on **Set**:

Top $\cong (\mathbb{F}, 2)$ -**Cat**

More generally:

T-Mon $\cong (\mathbb{T}, 2)$ -**Cat**, for \mathbb{T} "power-enriched"

A glance at properties: Hausdorffness, compactness...

Trivially for a relation r :

$$r : X \rightrightarrows Y \text{ is a map} \iff \begin{array}{ll} r \cdot r^\circ \leq 1_Y & r(x) \text{ uniquely defined} \\ 1_X \leq r^\circ \cdot r & r(x) \text{ defined everywhere} \end{array}$$

[Kamnitzer 1974]

$$\begin{array}{ll} a : TX \rightrightarrows X \text{ Hausdorff:} & a \cdot a^\circ \leq 1_X \quad \text{“}x \text{ has at most one pt of conv”} \\ \text{compact:} & 1_{TX} \leq a^\circ \cdot a \quad \text{“}x \text{ has at least one pt of conv”} \end{array}$$

If V is such that V -relations satisfying both inequalities are maps:

$(\mathbb{T}, V)\text{-Cat}_{\text{CompHaus}}$ is the Eilenberg-Moore category of \mathbb{T}

For $\mathbb{T} = \mathbb{U}, V = 2$:

[Manes 1969] **CompHaus** is monadic over **Set**.

A glance at properties: Hausdorffness, compactness...

Trivially for a relation r :

$$r : X \rightrightarrows Y \text{ is a map} \iff \begin{array}{ll} r \cdot r^\circ \leq 1_Y & r(x) \text{ uniquely defined} \\ 1_X \leq r^\circ \cdot r & r(x) \text{ defined everywhere} \end{array}$$

[Kamnitzer 1974]

$$\begin{array}{ll} a : TX \rightrightarrows X \text{ Hausdorff:} & a \cdot a^\circ \leq 1_X \quad \text{“}x \text{ has at most one pt of conv”} \\ \text{compact:} & 1_{TX} \leq a^\circ \cdot a \quad \text{“}x \text{ has at least one pt of conv”} \end{array}$$

If V is such that V -relations satisfying both inequalities are maps:

$(\mathbb{T}, V)\text{-Cat}_{\text{CompHaus}}$ is the Eilenberg-Moore category of \mathbb{T}

For $\mathbb{T} = \mathbb{U}, V = 2$:

[Manes 1969] **CompHaus** is monadic over **Set**,

A glance at properties: Hausdorffness, compactness...

Trivially for a relation r :

$$r : X \rightrightarrows Y \text{ is a map} \iff \begin{array}{ll} r \cdot r^\circ \leq 1_Y & r(x) \text{ uniquely defined} \\ 1_X \leq r^\circ \cdot r & r(x) \text{ defined everywhere} \end{array}$$

[Kamnitzer 1974]

$$\begin{array}{ll} a : TX \rightrightarrows X \text{ Hausdorff:} & a \cdot a^\circ \leq 1_X \quad \text{“}x \text{ has at most one pt of conv”} \\ \text{compact:} & 1_{TX} \leq a^\circ \cdot a \quad \text{“}x \text{ has at least one pt of conv”} \end{array}$$

If V is such that V -relations satisfying both inequalities are maps:

$(\mathbb{T}, V)\text{-Cat}_{\text{CompHaus}}$ is the Eilenberg-Moore category of \mathbb{T}

For $\mathbb{T} = \mathbb{U}, V = 2$:

[Manes 1969] **CompHaus** is monadic over **Set**,

A glance at properties: Hausdorffness, compactness...

Trivially for a relation r :

$$r : X \rightrightarrows Y \text{ is a map} \iff \begin{array}{ll} r \cdot r^\circ \leq 1_Y & r(x) \text{ uniquely defined} \\ 1_X \leq r^\circ \cdot r & r(x) \text{ defined everywhere} \end{array}$$

[Kamnitzer 1974]

$$\begin{array}{ll} a : TX \rightrightarrows X \text{ Hausdorff:} & a \cdot a^\circ \leq 1_X \quad \text{“}x \text{ has at most one pt of conv”} \\ \text{compact:} & 1_{TX} \leq a^\circ \cdot a \quad \text{“}x \text{ has at least one pt of conv”} \end{array}$$

If V is such that V -relations satisfying both inequalities are maps:

$(\mathbb{T}, V)\text{-Cat}_{\text{CompHaus}}$ is the Eilenberg-Moore category of \mathbb{T}

For $\mathbb{T} = \mathbb{U}, V = 2$:

[Manes 1969] **CompHaus** is monadic over **Set**.

... also being equationally defined in a lax environment

(\mathbb{T}, \mathbb{V}) -**Cat** :

$f : (X, a) \longrightarrow (Y, b)$ characterized by: $f \cdot a \leq b \cdot f$
 f (\mathbb{T}, \mathbb{V}) -**proper** defined by: $f \cdot a = b \cdot f$

Under mild conditions on \mathbb{T}, \mathbb{V} :

X Hausdorff $\iff X \longrightarrow X \times X$ proper

X compact $\iff X \longrightarrow 1$ proper

Other important equationally-defined map and object concepts:
open, local homeomorphism, regular, exponentiable, ...

[Hofmann-Seal-T (eds) 2014]: *Monoidal Topology*

... also being equationally defined in a lax environment

(\mathbb{T}, \mathbb{V}) -**Cat** :

$f : (X, a) \longrightarrow (Y, b)$ characterized by: $f \cdot a \leq b \cdot f$
 f (\mathbb{T}, \mathbb{V}) -proper defined by: $f \cdot a = b \cdot f$

Under mild conditions on \mathbb{T}, \mathbb{V} :

X Hausdorff $\iff X \longrightarrow X \times X$ proper

X compact $\iff X \longrightarrow 1$ proper

Other important equationally-defined map and object concepts:
open, local homeomorphism, regular, exponentiable, ...

[Hofmann-Seal-T (eds) 2014]: *Monoidal Topology*

Finally: Letting Hausdorff's target set N "float"

[Kopperman 1988]

Continuity space $(X, d: X \times X \rightarrow A, (A, +), P \subseteq A)$

"All topologies come from generalized metrics"

[Henriksen-Kopperman 1991]

Structure space $(X, d: X \times X \rightarrow A, A = 2^R)$

Transition to keeping " N " fixed: [Flagg 1997], [Flagg-Kopperman 1997]

V-continuity space: essentially V-**Cat**!

A more recent "floater": [Dikranjan-Zava 2017]

Coarse space (via ballean): $(X, P, B: X \times P \rightarrow 2^X)$

Rewrite B as $X \times X \rightarrow 2^P$: see structure space above!

Finally: Letting Hausdorff's target set N "float"

[Kopperman 1988]

Continuity space $(X, d: X \times X \rightarrow A, (A, +), P \subseteq A)$

"All topologies come from generalized metrics"

[Henriksen-Kopperman 1991]

Structure space $(X, d: X \times X \rightarrow A, A = 2^R)$

Transition to keeping " N " fixed: [Flagg 1997], [Flagg-Kopperman 1997]

V-continuity space: essentially V-Cat!

A more recent "floater": [Dikranjan-Zava 2017]

Coarse space (via ballean): $(X, P, B: X \times P \rightarrow 2^X)$

Rewrite B as $X \times X \rightarrow 2^P$: see structure space above!

Finally: Letting Hausdorff's target set N "float"

[Kopperman 1988]

Continuity space $(X, d: X \times X \rightarrow A, (A, +), P \subseteq A)$

"All topologies come from generalized metrics"

[Henriksen-Kopperman 1991]

Structure space $(X, d: X \times X \rightarrow A, A = 2^R)$

Transition to keeping " N " fixed: [Flagg 1997], [Flagg-Kopperman 1997]

V-continuity space: essentially V-**Cat!**

A more recent "floater": [Dikranjan-Zava 2017]

Coarse space (via ballean): $(X, P, B: X \times P \rightarrow 2^X)$

Rewrite B as $X \times X \rightarrow 2^P$: see structure space above!

Finally: Letting Hausdorff's target set N "float"

[Kopperman 1988]

Continuity space $(X, d: X \times X \rightarrow A, (A, +), P \subseteq A)$

"All topologies come from generalized metrics"

[Henriksen-Kopperman 1991]

Structure space $(X, d: X \times X \rightarrow A, A = 2^R)$

Transition to keeping " N " fixed: [Flagg 1997], [Flagg-Kopperman 1997]

V-continuity space: essentially V-**Cat**!

A more recent "floater": [Dikranjan-Zava 2017]

Coarse space (via ballean): $(X, P, B: X \times P \rightarrow 2^X)$

Rewrite B as $X \times X \rightarrow 2^P$: see structure space above!

Selected milestones leading to (\mathbb{T}, V) -Cat, and beyond

- Enriched category theory:
[Eilenberg-Kelly 1966], [Lawvere 1973], [Kelly 1982], ...
- Monad theory and its application to topology:
[Eilenberg-Moore 1965], [Kleisli 1965], [Manes 1969], [Barr 1970], ..
- Combining the two:
[Clementino-Hofmann 2003], [Clementino-T 2003],
[Clementino-Hofmann T 2003], [Seal 2005], [Hofmann 2007],
[Hofmann-Seal-T (eds) 2014], [T 2016], [Hofmann-Nora 2017], ...
- Resources from categorical topology:
[Brümmer 1971], [Penon 1972], [Herrlich 1974], [Manes 1974],
[Kamnitzer 1974], [Möbus 1981], [Wyler 1995], [Lowen 1997],
[Flagg 1997], [T 1999], [Höhle 2001], [Manes 2010], ...

Selected milestones leading to (\mathbb{T}, \mathbb{V}) -Cat, and beyond

- Enriched category theory:
[Eilenberg-Kelly 1966], [Lawvere 1973], [Kelly 1982], ...
- Monad theory and its application to topology:
[Eilenberg-Moore 1965], [Kleisli 1965], [Manes 1969], [Barr 1970], ..
- Combining the two:
[Clementino-Hofmann 2003], [Clementino-T 2003],
[Clementino-Hofmann T 2003], [Seal 2005], [Hofmann 2007],
[Hofmann-Seal-T (eds) 2014], [T 2016], [Hofmann-Nora 2017], ...
- Resources from categorical topology:
[Brümmer 1971], [Penon 1972], [Herrlich 1974], [Manes 1974],
[Kamnitzer 1974], [Möbus 1981], [Wyler 1995], [Lowen 1997],
[Flagg 1997], [T 1999], [Höhle 2001], [Manes 2010], ...

Selected milestones leading to (\mathbb{T}, \mathbb{V}) -Cat, and beyond

- Enriched category theory:
[Eilenberg-Kelly 1966], [Lawvere 1973], [Kelly 1982], ...
- Monad theory and its application to topology:
[Eilenberg-Moore 1965], [Kleisli 1965], [Manes 1969], [Barr 1970], ..
- Combining the two:
[Clementino-Hofmann 2003], [Clementino-T 2003],
[Clementino-Hofmann T 2003], [Seal 2005], [Hofmann 2007],
[Hofmann-Seal-T (eds) 2014], [T 2016], [Hofmann-Nora 2017], ...
- Resources from categorical topology:
[Brümmer 1971], [Penon 1972], [Herrlich 1974], [Manes 1974],
[Kamnitzer 1974], [Möbus 1981], [Wyler 1995], [Lowen 1997],
[Flagg 1997], [T 1999], [Höhle 2001], [Manes 2010], ...

Selected milestones leading to (\mathbb{T}, V) -Cat, and beyond

- Enriched category theory:
[Eilenberg-Kelly 1966], [Lawvere 1973], [Kelly 1982], ...
- Monad theory and its application to topology:
[Eilenberg-Moore 1965], [Kleisli 1965], [Manes 1969], [Barr 1970], ..
- Combining the two:
[Clementino-Hofmann 2003], [Clementino-T 2003],
[Clementino-Hofmann T 2003], [Seal 2005], [Hofmann 2007],
[Hofmann-Seal-T (eds) 2014], [T 2016], [Hofmann-Nora 2017], ...
- Resources from categorical topology:
[Brümmer 1971], [Penon 1972], [Herrlich 1974], [Manes 1974],
[Kamnitzer 1974], [Möbus 1981], [Wyler 1995], [Lowen 1997],
[Flagg 1997], [T 1999], [Höhle 2001], [Manes 2010], ...

Your questions, comments?

THANK YOU !