

1995

Positive solutions for higher order ordinary differential equations

Paul W. Eloë

University of Dayton, peloe1@udayton.edu

Johnny Henderson

Baylor University

Follow this and additional works at: https://ecommons.udayton.edu/mth_fac_pub



Part of the [Mathematics Commons](#)

eCommons Citation

Eloë, Paul W. and Henderson, Johnny, "Positive solutions for higher order ordinary differential equations" (1995). *Mathematics Faculty Publications*. 98.

https://ecommons.udayton.edu/mth_fac_pub/98

This Article is brought to you for free and open access by the Department of Mathematics at eCommons. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of eCommons. For more information, please contact frice1@udayton.edu, mschlangen1@udayton.edu.

POSITIVE SOLUTIONS FOR HIGHER ORDER ORDINARY DIFFERENTIAL EQUATIONS

PAUL W. ELOE & JOHNNY HENDERSON

ABSTRACT. Solutions that are positive with respect to a cone are obtained for the boundary value problem, $u^{(n)} + a(t)f(u) = 0$, $u^{(i)}(0) = u^{(n-2)}(1) = 0$, $0 \leq i \leq n-2$, in the cases that f is either superlinear or sublinear. The methods involve application of a fixed point theorem for operators on a cone.

1. INTRODUCTION

We are concerned with the existence of solutions for the two-point boundary value problem,

$$u^{(n)} + a(t)f(u) = 0, \quad 0 < t < 1, \quad (1)$$

$$u^{(i)}(0) = u^{(n-2)}(1) = 0, \quad 0 \leq i \leq n-2, \quad (2)$$

where

(A) $f : [0, \infty) \rightarrow [0, \infty)$ is continuous, and

(B) $a : [0, 1] \rightarrow [0, \infty)$ is continuous and does not vanish identically on any subinterval.

We remark that, if $u(t)$ is a nonnegative solution of (1), (2), then $u^{(n-2)}(t)$ is concave on $[0, 1]$.

Specifically, our aim is to extend the work of Erbe and Wang [10] to obtain solutions of (1), (2), that are positive with respect to a cone, in the cases when, either (i) f is superlinear, or (ii) f is sublinear; that is, in the respective cases when, either (i) $f_0 = 0$ and $f_\infty = \infty$, or (ii) $f_0 = \infty$ and $f_\infty = 0$, where

$$f_0 = \lim_{x \rightarrow 0^+} \frac{f(x)}{x} \quad \text{and} \quad f_\infty = \lim_{x \rightarrow \infty} \frac{f(x)}{x}.$$

In the case that $n = 2$, the boundary value problem (1), (2) arises in applications involving nonlinear elliptic problems in annular regions; see [1], [2], [12], [19]. Applications of (1), (2) can also be made to singular boundary value problems as in [3], [6], [8], [13], [16], [18], as well as to extremal point characterizations for

1991 *Mathematics Subject Classification.* 34B15.

Key words and phrases. Boundary value problems, positive solutions, superlinear and sublinear, operators on a cone.

©1995 Southwest Texas State University and University of North Texas.

Submitted on December 4, 1994.

boundary value problems in [9], [14], [17]. In these applications, frequently, only solutions that are positive are useful. The results herein are also somewhat related to those obtained in [5] and [11].

Our arguments for establishing the existence of solutions of (1), (2) involve concavity properties of solutions that are used in defining a cone on which a positive integral operator is defined. A fixed point theorem due to Krasnosel'skii [15] is applied to yield a positive solution of (1), (2).

In Section 2, we present some properties of a Green's function which will be used in defining the positive operator. We also state the fixed point theorem from [15]. In Section 3, we provide an appropriate Banach space and cone in order to apply the fixed point theorem yielding solutions of (1), (2) in both the superlinear and sublinear cases.

2. SOME PRELIMINARIES

In this section, we state a theorem due to Krasnosel'skii, an application of which will yield in the next section a positive solution of (1), (2). The mapping to which we apply this fixed point theorem will include an integral whose kernel, $G(t, s)$, is the Green's function for

$$\begin{aligned} -y^{(n)} &= 0, \\ y^{(i)}(0) = y^{(n-2)}(1) &= 0, \quad 0 \leq i \leq n-2. \end{aligned} \tag{3}$$

Eloe [7] has shown that, for $0 \leq i \leq n-2$,

$$\frac{\partial^i}{\partial t^i} G(t, s) > 0 \text{ on } (0, 1) \times (0, 1), \tag{4}$$

as well as the fact that the function

$$K(t, s) = \frac{\partial^{n-2}}{\partial t^{n-2}} G(t, s) \tag{5}$$

is the Green's function for

$$\begin{aligned} -y'' &= 0, \\ y(0) = y(1) &= 0. \end{aligned} \tag{6}$$

We note that

$$K(t, s) = \begin{cases} t(1-s), & 0 \leq t < s \leq 1, \\ s(1-t), & 0 \leq s < t \leq 1, \end{cases} \tag{7}$$

from which it is straightforward that

$$K(t, s) \leq K(s, s), \quad 0 \leq t, s \leq 1, \tag{8}$$

and a nice argument in [10] shows that

$$K(t, s) \geq \frac{1}{4} K(s, s), \quad \frac{1}{4} \leq t \leq \frac{3}{4}, \quad 0 \leq s \leq 1. \tag{9}$$

The existence of solutions of (1), (2) is based on an application of the following fixed point theorem [15].

Theorem 1. Let \mathcal{B} be a Banach space, and let $\mathcal{P} \subset \mathcal{B}$ be a cone in \mathcal{B} . Assume Ω_1, Ω_2 are open subsets of \mathcal{B} with $0 \in \Omega_1 \subset \bar{\Omega}_1 \subset \Omega_2$, and let

$$T : \mathcal{P} \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow \mathcal{P}$$

be a completely continuous operator such that, either

- (i) $\|Tu\| \leq \|u\|$, $u \in \mathcal{P} \cap \partial\Omega_1$, and $\|Tu\| \geq \|u\|$, $u \in \mathcal{P} \cap \partial\Omega_2$, or
- (ii) $\|Tu\| \geq \|u\|$, $u \in \mathcal{P} \cap \partial\Omega_1$, and $\|Tu\| \leq \|u\|$, $u \in \mathcal{P} \cap \partial\Omega_2$.

Then T has a fixed point in $\mathcal{P} \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

3. EXISTENCE OF SOLUTIONS

We are now ready to apply Theorem 1. We remark that $u(t)$ is a solution of (1), (2) if, and only if,

$$u(t) = \int_0^1 G(t,s)a(s)f(u(s))ds, \quad 0 \leq t \leq 1.$$

For our construction, we let

$$\mathcal{B} = \{x \in C^{(n-2)}[0,1] \mid x^{(i)}(0) = 0, \quad 0 \leq i \leq n-3\},$$

with norm, $\|x\| = |x^{(n-2)}|_\infty$, where $|\cdot|_\infty$ denotes the supremum norm on $[0,1]$. Then $(\mathcal{B}, \|\cdot\|)$ is a Banach space.

Remark 1. We note that, for each $x \in \mathcal{B}$,

$$|x^{(i)}|_\infty \leq \|x\|, \quad 0 \leq i \leq n-2. \quad (10)$$

We will seek solutions of (1), (2) which lie in a cone, \mathcal{P} , defined by

$$\mathcal{P} = \{x \in \mathcal{B} \mid x^{(n-2)}(t) \geq 0 \text{ on } [0,1], \text{ and } \min_{\frac{1}{4} \leq t \leq \frac{3}{4}} x^{(n-2)}(t) \geq \frac{1}{4}\|x\|\}.$$

Remark 2. We note here that, if $x \in \mathcal{P}$, then $x^{(i)}(t) \geq 0$ on $[0,1]$ and

$$x^{(i)}(t) \geq \frac{1}{4}\|x\| \frac{(t - \frac{1}{4})^{n-i-2}}{(n-i-2)!}$$

on $[\frac{1}{4}, \frac{3}{4}]$, $0 \leq i \leq n-2$. As a consequence

$$x^{(i)}(t) \geq \frac{1}{(n-i-2)!4^{n-i-1}}\|x\|$$

on $[\frac{1}{2}, \frac{3}{4}]$, $0 \leq i \leq n-2$.

Theorem 2. Assume that conditions (A) and (B) are satisfied. If, either

- (i) $f_0 = 0$ and $f_\infty = \infty$ (i.e., f is superlinear), or
- (ii) $f_0 = \infty$ and $f_\infty = 0$ (i.e., f is sublinear),

then (1), (2) has at least one solution in \mathcal{P} .

Proof. We begin by defining an integral operator $T : \mathcal{P} \rightarrow \mathcal{B}$ by

$$Tu(t) = \int_0^1 G(t,s)a(s)f(u(s))ds, \quad u \in \mathcal{P}, \quad (11)$$

and we seek a fixed point of T in the cone \mathcal{P} for the respective cases of f superlinear and f sublinear.

Before dealing with these cases, we make a few observations. First, if $u \in \mathcal{P}$, it follows from (8) that

$$\begin{aligned} (Tu)^{(n-2)}(t) &= \int_0^1 \frac{\partial^{n-2}}{\partial t^{n-2}} G(t, s) a(s) f(u(s)) ds \\ &= \int_0^1 K(t, s) a(s) f(u(s)) ds \\ &\leq \int_0^1 K(s, s) a(s) f(u(s)) ds, \end{aligned}$$

so that

$$\|Tu\| = |(Tu)^{(n-2)}|_\infty \leq \int_0^1 K(s, s) a(s) f(u(s)) ds.$$

In fact,

$$\|Tu\| = \int_0^1 K(s, s) a(s) f(u(s)) ds. \quad (12)$$

Next, if $u \in \mathcal{P}$, it follows from (9) and (12) that

$$\begin{aligned} \min_{\frac{1}{4} \leq t \leq \frac{3}{4}} (Tu)^{(n-2)}(t) &= \min_{\frac{1}{4} \leq t \leq \frac{3}{4}} \int_0^1 K(t, s) a(s) f(u(s)) ds \\ &\geq \frac{1}{4} \int_0^1 K(s, s) a(s) f(u(s)) ds \\ &\geq \frac{1}{4} \|Tu\|. \end{aligned}$$

Moreover, properties of $G(t, s)$ give that $(Tu)^{(n-2)}(t) \geq 0$, so that $Tu \in \mathcal{P}$, and in particular $T : \mathcal{P} \rightarrow \mathcal{P}$. Also, the standard arguments yield that T is completely continuous.

We now turn to the cases of the theorem.

- (i) Assume $f_0 = 0$ and $f_\infty = \infty$. First, dealing with $f_0 = 0$, there exist $\eta > 0$ and $H_1 > 0$ such that $f(x) \leq \eta x$, for $0 < x \leq H_1$, and

$$\eta \int_0^1 K(s, s) a(s) ds \leq 1.$$

So, if we choose $u \in \mathcal{P}$ with $\|u\| = H_1$, and if we recall from Remark 1 that $|u|_\infty \leq \|u\|$, we have from (8),

$$\begin{aligned}
(Tu)^{(n-2)}(t) &= \int_0^1 K(t,s)a(s)f(u(s)) ds \\
&\leq \int_0^1 K(s,s)a(s)f(u(s)) ds \\
&\leq \int_0^1 K(s,s)a(s)\eta u(s) ds \\
&\leq \eta \int_0^1 K(s,s)a(s) ds \|u\| \\
&\leq \|u\|, \quad 0 \leq t \leq 1.
\end{aligned}$$

As a consequence $\|Tu\| = |(Tu)^{(n-2)}(t)|_\infty \leq \|u\|$. Thus, if we set

$$\Omega_1 = \{x \in \mathcal{B} \mid \|x\| < H_1\},$$

then

$$\|Tu\| \leq \|u\|, \text{ for } u \in \mathcal{P} \cap \partial\Omega_1. \quad (13)$$

Next, dealing with $f_\infty = \infty$, there exist $\lambda > 0$ and $\bar{H}_2 > 0$ such that $f(x) \geq \lambda x$, for $x \geq \bar{H}_2$, and

$$\frac{\lambda}{(n-2)!4^{n-1}} \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right)a(s) ds \geq 1.$$

Now, let $H_2 = \max\{2H_1, (n-2)!4^{n-1}\bar{H}_2\}$ and set

$$\Omega_2 = \{x \in \mathcal{B} \mid \|x\| < H_2\}.$$

So, if $u \in \mathcal{P}$ with $\|u\| = H_2$, and if we recall from Remark 2 that $u(t) \geq \frac{1}{(n-2)!4^{n-1}}\|u\| \geq \bar{H}_2$ on $[\frac{1}{2}, \frac{3}{4}]$, we have

$$\begin{aligned}
(Tu)^{(n-2)}\left(\frac{1}{2}\right) &= \int_0^1 K\left(\frac{1}{2}, s\right)a(s)f(u(s)) ds \\
&\geq \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right)a(s)\lambda u(s) ds \\
&\geq \lambda \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right)a(s) \frac{1}{(n-2)!4^{n-1}} \|u\| ds \\
&= \frac{\lambda}{(n-2)!4^{n-1}} \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right)a(s) ds \|u\| \\
&\geq \|u\|,
\end{aligned}$$

so that $\|Tu\| \geq \|u\|$. Consequently,

$$\|Tu\| \geq \|u\|, \text{ for } u \in \mathcal{P} \cap \partial\Omega_2. \quad (14)$$

Therefore, by part (i) of Theorem 1 applied to (13) and (14), T has a fixed point $u(t) \in \mathcal{P} \cap (\Omega_2 \setminus \Omega_1)$ such that $H_1 \leq \|u\| \leq H_2$, and as such, $u(t)$ is a desired solution of (1), (2). (We remark that the arguments carry through, if we had set $H_2 = \max\{H_1, (n-2)!4^{n-1}\bar{H}_2\}$ and if $H_2 = H_1$, then there

is a solution $u \in \mathcal{P}$ with $\|u\| = H_1$.) This completes the case when f is superlinear.

- (ii) Now, assume $f_0 = \infty$ and $f_\infty = 0$. Dealing with $f_0 = \infty$, there exist $\bar{\eta} > 0$ and $J_1 > 0$ such that $f(x) \geq \bar{\eta}x$, for $0 < x \leq J_1$, and

$$\frac{\bar{\eta}}{(n-2)!4^{n-1}} \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right) a(s) ds \geq 1.$$

This time, we choose $u \in \mathcal{P}$ with $\|u\| = J_1$. Since $|u|_\infty \leq \|u\| = J_1$, we have $f(u(s)) \geq \bar{\eta}u(s)$, $0 \leq s \leq 1$. Also, we know $u(s) \geq \frac{1}{(n-2)!4^{n-1}}\|u\|$, $\frac{1}{2} \leq s \leq \frac{3}{4}$. Thus,

$$\begin{aligned} (Tu)^{(n-2)}\left(\frac{1}{2}\right) &= \int_0^1 K\left(\frac{1}{2}, s\right) a(s) f(u(s)) ds \\ &\geq \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right) a(s) \bar{\eta}u(s) ds \\ &\geq \frac{\bar{\eta}}{(n-2)!4^{n-1}} \int_{\frac{1}{2}}^{\frac{3}{4}} K\left(\frac{1}{2}, s\right) a(s) ds \|u\| \\ &\geq \|u\|, \end{aligned}$$

and in particular, $\|Tu\| \geq \|u\|$. Setting

$$\Omega_1 = \{x \in \mathcal{B} \mid \|x\| < J_1\},$$

we conclude

$$\|Tu\| \geq \|u\|, \text{ for } u \in \mathcal{P} \cap \partial\Omega_1. \quad (15)$$

For the final part of this case, we deal with $f_\infty = 0$. There exist $\bar{\lambda} > 0$ and $\bar{J}_2 > 0$ such that, $f(x) \leq \bar{\lambda}x$, for $x \geq \bar{J}_2$, and

$$\bar{\lambda} \int_0^1 K(s, s) a(s) ds \leq 1.$$

There are two further sub-cases to be considered:

- (I) We suppose first that f is bounded. Then, there exists $N > 0$ such that $f(x) \leq N$, for all $0 < x < \infty$. Let $J_2 = \max\{2J_1, N \int_0^1 K(s, s) a(s) ds\}$. Then, for $u \in \mathcal{P}$ with $\|u\| = J_2$, since $|u|_\infty \leq \|u\|$ and $K(t, s) \leq K(s, s)$, $0 \leq s, t \leq 1$, we have

$$\begin{aligned} (Tu)^{(n-2)}(t) &= \int_0^1 K(t, s) a(s) f(u(s)) ds \\ &\leq N \int_0^1 K(s, s) a(s) ds \\ &\leq J_2 \\ &= \|u\|, \quad 0 \leq t \leq 1. \end{aligned}$$

Consequently, $\|Tu\| \leq \|u\|$.

- (II) For the second sub-case, suppose that f is unbounded. Then, there exists $J_2 > \max\{2J_1, \bar{J}_2\}$ such that $f(x) \leq f(J_2)$, for $0 < x \leq J_2$. We now choose $u \in \mathcal{P}$ with $\|u\| = J_2$. Again, recalling $|u|_\infty \leq \|u\|$ and $K(t, s) \leq K(s, s)$ leads to

$$\begin{aligned}
(Tu)^{(n-2)}(t) &= \int_0^1 K(t,s)a(s)f(u(s)) ds \\
&\leq \int_0^1 K(s,s)a(s)f(J_2) ds \\
&\leq \bar{\lambda} \int_0^1 K(s,s)a(s) ds J_2 \\
&\leq \|u\|, \quad 0 \leq t \leq 1.
\end{aligned}$$

Thus, $\|Tu\| \leq \|u\|$.

We conclude from each sub-case, (I) and (II), if we set

$$\Omega_2 = \{x \in \mathcal{B} \mid \|x\| < J_2\},$$

then

$$\|Tu\| \leq \|u\|, \text{ for } u \in \mathcal{P} \cap \partial\Omega_2. \quad (16)$$

Therefore, by part (ii) of Theorem 1 applied to (15) and (16), T has a fixed point $u(t) \in \mathcal{P} \cap (\Omega_2 \setminus \Omega_1)$ such that $J_1 \leq \|u\| \leq J_2$, and $u(t)$ is a sought solution of (1), (2). This completes the argument for the case of f sublinear.

The proof is complete. \square

REFERENCES

- [1] C. Bandle, C. V. Coffman and M. Marcus, Nonlinear elliptic problems in annular domains, *J. Diff. Eqns.* **69** (1987), 322-345.
- [2] C. Bandle and M. K. Kwong, Semilinear elliptic problems in annular domains, *J. Appl. Math. Phys.* **40** (1989), 245-257.
- [3] F. Bernis, On some nonlinear singular boundary value problems of higher order, *Nonlin. Anal.*, in press.
- [4] A. Callegari and A. Nachman, A nonlinear singular boundary value problem in the theory of pseudoplastic fluids, *SIAM J. Appl. Math.* **38** (1980), 275-281.
- [5] C. J. Chyan, Positive solution of a system of higher order boundary value problems, *PanAm. Math. J.*, in press.
- [6] C. J. Chyan and J. Henderson, Positive solutions for singular higher order nonlinear equations, *Diff. Eqns. Dyn. Sys.* **2** (1994), 153-160.
- [7] P. W. Eloe, Sign properties of Green's functions for two classes of boundary value problems, *Canad. Math. Bull.* **30** (1987), 28-35.
- [8] P. W. Eloe and J. Henderson, Singular nonlinear boundary value problems for higher order ordinary differential equations, *Nonlin. Anal.* **17** (1991), 1-10.
- [9] P. W. Eloe and J. Henderson, Focal points and comparison theorems for a class of two point boundary value problems, *J. Diff. Eqns.* **103** (1993), 375-386.
- [10] L. H. Erbe and H. Wang, On the existence of positive solutions of ordinary differential equations, *Proc. Amer. Math. Soc.* **120** (1994), 743-748.
- [11] A. M. Fink, J. A. Gatica and G. E. Hernandez, Eigenvalues of generalized Gel'fand models, *Nonlin. Anal.* **20** (1993), 1453-1468.
- [12] X. Garaizar, Existence of positive radial solutions for semilinear elliptic problems in the annulus, *J. Diff. Eqns.* **70** (1987), 69-72.
- [13] J. A. Gatica, V. Oliker and P. Waltman, Singular nonlinear boundary value problems for second-order ordinary differential equations, *J. Diff. Eqns.* **79** (1989), 62-78.
- [14] D. Hankerson and J. Henderson, Positive solutions and extremal points for differential equations, *Appl. Anal.* **39** (1990), 193-207.
- [15] M. A. Krasnosel'skii, "Positive solutions of operator equations", Noordhoff, Groningen, 1964.
- [16] C. D. Luning and W. L. Perry, Positive solutions of negative exponent generalized Emden-Fowler boundary value problems, *SIAM J. Appl. Math.* **12** (1981), 874-879.

- [17] K. Schmitt and H. L. Smith, Positive solutions and conjugate points for systems of differential equations, *Nonlin. Anal.* **2** (1978), 93-105.
- [18] S. Taliaferro, A nonlinear singular boundary value problem, *Nonlin. Anal.* **3** (1979), 897-904.
- [19] H. Wang, On the existence of positive solutions for semilinear elliptic equations in the annulus, *J. Diff. Eqns.* **109** (1994), 1-7.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DAYTON, DAYTON, OH 45469-2316 USA
E-mail address: `eloe@udavxb.oca.udayton.edu`

DISCRETE AND STATISTICAL SCIENCES, AUBURN UNIVERSITY, AUBURN, AL 36849-5307 USA
E-mail address: `hendez2@mail.auburn.edu`