

2014

# Existence and comparison of smallest eigenvalues for a fractional boundary-value problem

Paul W. Eloë

*University of Dayton*, [peloe1@udayton.edu](mailto:peloe1@udayton.edu)

Jeffrey T. Neugebauer

*Eastern Kentucky University*

Follow this and additional works at: [https://ecommons.udayton.edu/mth\\_fac\\_pub](https://ecommons.udayton.edu/mth_fac_pub)



Part of the [Mathematics Commons](#)

---

## eCommons Citation

Eloë, Paul W. and Neugebauer, Jeffrey T., "Existence and comparison of smallest eigenvalues for a fractional boundary-value problem" (2014). *Mathematics Faculty Publications*. 102.

[https://ecommons.udayton.edu/mth\\_fac\\_pub/102](https://ecommons.udayton.edu/mth_fac_pub/102)

This Article is brought to you for free and open access by the Department of Mathematics at eCommons. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of eCommons. For more information, please contact [frice1@udayton.edu](mailto:frice1@udayton.edu), [mschlangen1@udayton.edu](mailto:mschlangen1@udayton.edu).

**EXISTENCE AND COMPARISON OF SMALLEST  
EIGENVALUES FOR A FRACTIONAL  
BOUNDARY-VALUE PROBLEM**

PAUL W. ELOE, JEFFREY T. NEUGEBAUER

ABSTRACT. The theory of  $u_0$ -positive operators with respect to a cone in a Banach space is applied to the fractional linear differential equations

$$D_{0+}^{\alpha}u + \lambda_1 p(t)u = 0 \quad \text{and} \quad D_{0+}^{\alpha}u + \lambda_2 q(t)u = 0,$$

$0 < t < 1$ , with each satisfying the boundary conditions  $u(0) = u(1) = 0$ . The existence of smallest positive eigenvalues is established, and a comparison theorem for smallest positive eigenvalues is obtained.

1. INTRODUCTION

We consider the eigenvalue problems

$$D_{0+}^{\alpha}u + \lambda_1 p(t)u = 0, \quad 0 < t < 1, \tag{1.1}$$

$$D_{0+}^{\alpha}u + \lambda_2 q(t)u = 0, \quad 0 < t < 1, \tag{1.2}$$

satisfying the boundary conditions

$$u(0) = u(1) = 0, \tag{1.3}$$

where  $1 < \alpha \leq 2$  is a real number,  $D_{0+}^{\alpha}$  is the standard Riemann-Liouville derivative, and  $p(t)$  and  $q(t)$  are continuous nonnegative functions on  $[0, 1]$ , where neither  $p(t)$  nor  $q(t)$  vanishes identically on any nondegenerate compact subinterval of  $[0, 1]$ .

The Krein Rutman theory [14] has been employed extensively to establish the existence of and compare smallest eigenvalues of boundary value problems for differential equations, difference equations, and dynamic equations on time scales. For some examples, see [4, 5, 7, 8, 9, 11, 12, 16, 18] and the references therein. A standard approach to show the existence of smallest eigenvalues is to apply the theory of  $u_0$ -positive operators [15]. Operators are defined whose eigenvalues are reciprocals of the eigenvalues of the original boundary value problems. These operators are constructed by using the corresponding Green's function; the  $u_0$ -positivity of these operators are obtained by showing the operator maps nonzero elements of a cone into the interior of that cone. Sign properties of the Green's function are employed to map the cone into the cone and higher order derivatives of the Green's

---

2000 *Mathematics Subject Classification.* 26A33.

*Key words and phrases.* Fractional boundary value problem; smallest eigenvalues;  $u_0$ -positive operator.

©2014 Texas State University - San Marcos.

Submitted August 22, 2013. Published February 10, 2014.

functions are employed to map elements to the interior of the cone. The theory of  $u_0$ -positivity, as developed by Krasnosel'skii [15], gives the existence of largest eigenvalues of the operator, with the corresponding eigenfunction existing in a cone.

In this article, we apply the standard approach, described above, to a boundary value problem for a fractional differential equation. We are not aware of any previous application of  $u_0$ -positive operators to fractional differential equations. Fixed point theory is now commonly applied to boundary value problems for fractional equations; see, for example, the bibliography found in [1]. In many of these applications, the common Banach space to employ is  $C[0, 1]$ ; this space is not appropriate for applications of  $u_0$ -positivity to (1.1), (1.2) or (1.1), (1.3), since the corresponding Green's function,  $G(t, s)$ , has unbounded slope at  $t = 0$ . The primary contribution of this article then is to consider an appropriate Banach space and cone, with nonempty interior, so that theory of  $u_0$ -positive operators does apply. The motivation for the Banach space used here is found in [6, Theorem 2.5] or [17, Theorem 3.1]. The particular approach to construct the Banach space and cone is modeled after [16]. For other work on eigenvalue problems of fractional differential equations, see [2, 10, 19, 20].

In Section 2, we state the preliminary definitions and theorems. In Section 3, we define the appropriate Banach space and establish the existence of and compare smallest eigenvalues of (1.1), (1.2) and (1.1), (1.3).

## 2. PRELIMINARY DEFINITIONS AND THEOREMS

**Definition 2.1.** Let  $1 < \alpha \leq 2$ . The  $\alpha$ -th Riemann-Liouville fractional derivative of the function  $u : [0, 1] \rightarrow \mathbb{R}$ , denoted  $D_{0+}^\alpha u$ , is defined as

$$D_{0+}^\alpha u(t) = \frac{1}{\Gamma(2-\alpha)} \frac{d^2}{dt^2} \int_0^t (t-s)^{2-\alpha-1} u(s) ds,$$

provided the right-hand side exists.

**Definition 2.2.** Let  $\mathcal{B}$  be a Banach space over  $\mathbb{R}$ . A closed nonempty subset  $\mathcal{P}$  of  $\mathcal{B}$  is said to be a cone provided

- (i)  $\alpha u + \beta v \in \mathcal{P}$ , for all  $u, v \in \mathcal{P}$  and all  $\alpha, \beta \geq 0$ , and
- (ii)  $u \in \mathcal{P}$  and  $-u \in \mathcal{P}$  implies  $u = 0$ .

**Definition 2.3.** A cone  $\mathcal{P}$  is solid if the interior,  $\mathcal{P}^\circ$ , of  $\mathcal{P}$ , is nonempty. A cone  $\mathcal{P}$  is reproducing if  $\mathcal{B} = \mathcal{P} - \mathcal{P}$ ; i.e., given  $w \in \mathcal{B}$ , there exist  $u, v \in \mathcal{P}$  such that  $w = u - v$ .

**Remark 2.4.** Krasnosel'skii [15] showed that every solid cone is reproducing.

Cones give rise to partial orders on Banach spaces and to partial orders on bounded linear operators on Banach spaces in a natural way.

**Definition 2.5.** Let  $\mathcal{P}$  be a cone in a real Banach space  $\mathcal{B}$ . If  $u, v \in \mathcal{B}$ , we say  $u \leq v$  with respect to  $\mathcal{P}$  if  $v - u \in \mathcal{P}$ . If both  $M, N : \mathcal{B} \rightarrow \mathcal{B}$  are bounded linear operators, we say  $M \leq N$  with respect to  $\mathcal{P}$  if  $Mu \leq Nu$  for all  $u \in \mathcal{P}$ .

**Definition 2.6.** A bounded linear operator  $M : \mathcal{B} \rightarrow \mathcal{B}$  is  $u_0$ -positive with respect to  $\mathcal{P}$  if there exists  $u_0 \in \mathcal{P} \setminus \{0\}$  such that for each  $u \in \mathcal{P} \setminus \{0\}$ , there exist  $k_1(u) > 0$  and  $k_2(u) > 0$  such that  $k_1 u_0 \leq Mu \leq k_2 u_0$  with respect to  $\mathcal{P}$ .

The following two results are fundamental to our comparison results and are attributed to Krasnosel'skii [15]. The proof of Theorem 2.7 can be found in Krasnosel'skii's book [15], and the proof of Theorem 2.8 is provided by Keener and Travis [13] as an extension of Krasnosel'skii's results.

**Theorem 2.7.** *Let  $\mathcal{B}$  be a real Banach space and let  $\mathcal{P} \subset \mathcal{B}$  be a reproducing cone. Let  $L : \mathcal{B} \rightarrow \mathcal{B}$  be a compact,  $u_0$ -positive, linear operator. Then  $L$  has an essentially unique eigenvector in  $\mathcal{P}$ , and the corresponding eigenvalue is simple, positive, and larger than the absolute value of any other eigenvalue.*

**Theorem 2.8.** *Let  $\mathcal{B}$  be a real Banach space and  $\mathcal{P} \subset \mathcal{B}$  be a cone. Let both  $M, N : \mathcal{B} \rightarrow \mathcal{B}$  be bounded, linear operators and assume that at least one of the operators is  $u_0$ -positive. If  $M \leq N$ ,  $Mu_1 \geq \lambda_1 u_1$  for some  $u_1 \in \mathcal{P}$  and some  $\lambda_1 > 0$ , and  $Nu_2 \leq \lambda_2 u_2$  for some  $u_2 \in \mathcal{P}$  and some  $\lambda_2 > 0$ , then  $\lambda_1 \leq \lambda_2$ . Furthermore,  $\lambda_1 = \lambda_2$  implies  $u_1$  is a scalar multiple of  $u_2$ .*

### 3. COMPARISON OF SMALLEST EIGENVALUES

In [3], Bai and Lu showed the Green's function for  $-D_{0+}^\alpha u(t) = 0$  satisfying (1.3) is

$$G(t, s) = \begin{cases} \frac{[t(1-s)]^{\alpha-1} - (t-s)^{\alpha-1}}{\Gamma(\alpha)}, & 0 \leq s \leq t \leq 1, \\ \frac{[t(1-s)]^{\alpha-1}}{\Gamma(\alpha)}, & 0 \leq t \leq s \leq 1. \end{cases} \quad (3.1)$$

Define the Banach Space

$$\mathcal{B} = \{u : u = t^{\alpha-1}v, v \in C^1[0, 1], v(1) = 0\},$$

with the norm

$$\|u\| = |v'|_0,$$

where  $|v'|_0 = \sup_{t \in [0, 1]} |v'(t)|$  denotes the usual supremum norm.

Note that for  $v \in C^1[0, 1]$ ,  $v(1) = 0$ ,  $0 \leq t \leq 1$ ,

$$|v(t)| = |v(t) - v(1)| = \left| \int_1^t v'(s) ds \right| \leq (1-t)|v'| \leq \|u\|.$$

Therefore,  $|v|_0 \leq \|u\| = |v'|_0$  and

$$|u|_0 = |t^{\alpha-1}v|_0 \leq t^{\alpha-1}\|u\|,$$

implies  $|u|_0 \leq \|u\|$ .

Define the linear operators

$$Mu(t) = \int_0^1 G(t, s)p(s)u(s)ds \quad (3.2)$$

and

$$Nu(t) = \int_0^1 G(t, s)q(s)u(s) ds. \quad (3.3)$$

**Theorem 3.1.** *The operators  $M, N : \mathcal{B} \rightarrow \mathcal{B}$  are compact linear operators.*

*Proof.* We first show  $M : \mathcal{B} \rightarrow \mathcal{B}$ . Let  $u \in \mathcal{B}$ . So there is a  $v \in C^1[0, 1]$  such that  $u = t^{\alpha-1}v$ . Since  $v \in C^1[0, 1]$  and  $p \in C[0, 1]$ , let  $L = |v|_0$  and let  $P = |p|_0$ . Now

$$Mu(t) = \int_0^1 \frac{t^{\alpha-1}(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds - \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds$$

$$= t^{\alpha-1} \left( \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds - t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds \right).$$

Define

$$g(t) = \begin{cases} 0, & t = 0, \\ t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds, & 0 < t \leq 1. \end{cases}$$

First, note  $g \in C^1(0, 1]$ . Now

$$\begin{aligned} |g(t)| &= \left| t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds \right| \\ &= \left| t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)s^{\alpha-1}v(s)ds \right| \\ &\leq PLt^{1-\alpha} \int_0^t (t-s)^{\alpha-1}s^{\alpha-1}ds \\ &\leq PLt^{1-\alpha}t^{\alpha-1} \int_0^t (t-s)^{\alpha-1}ds \\ &= \frac{PLt^\alpha}{\alpha}, \end{aligned}$$

where  $\frac{PL}{\alpha} \geq 0$ . So  $\lim_{t \rightarrow 0^+} g(t) = g(0) = 0$  and  $g \in C[0, 1]$ .

Also, for  $t > 0$ ,

$$\begin{aligned} |g'(t)| &= \left| (1-\alpha)t^{-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds \right. \\ &\quad \left. + (\alpha-1)t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha)} p(s)u(s)ds \right| \\ &\leq \left| (1-\alpha)t^{-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)s^{\alpha-1}v(s)ds \right| \\ &\quad + \left| t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha-1)} p(s)s^{\alpha-1}v(s)ds \right| \\ &\leq (\alpha-1)PLt^{-\alpha}t^{\alpha-1} \int_0^t (t-s)^{\alpha-1}ds + PLt^{1-\alpha}t^{\alpha-1} \int_0^t (t-s)^{\alpha-2}ds \\ &= \left( \frac{\alpha-1}{\alpha} + \frac{1}{\alpha-1} \right) PLt^{\alpha-1}. \end{aligned}$$

So,  $\lim_{t \rightarrow 0^+} g'(t) = 0$ . Moreover, using the definition of derivative and L'Hospital's rule,

$$g'(0) = \lim_{t \rightarrow 0^+} \frac{g(t) - g(0)}{t} = \lim_{t \rightarrow 0^+} \frac{g(t)}{t} = \lim_{t \rightarrow 0^+} g'(t) = 0,$$

and so  $g' \in C[0, 1]$ .

Now set

$$\hat{v}(t) = \left( \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds - t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds \right).$$

It is an easy calculation to verify that  $\hat{v}(1) = 0$ . Thus  $Mu \in \mathcal{B}$ . So  $M : \mathcal{B} \rightarrow \mathcal{B}$ . The proof that  $N : \mathcal{B} \rightarrow \mathcal{B}$  is similar.

We now show that  $M : \mathcal{B} \rightarrow \mathcal{B}$  is a compact operator. Let  $L > 0$  and consider

$$K = \{u \in \mathcal{B} : \|u\| \leq L\}$$

or more appropriately consider

$$\hat{K} = \{v \in C^1[0, 1] : v(1) = 0, |v'|_0 \leq L\}.$$

Define

$$\hat{M}(v)(t) = \left( \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s) s^{\alpha-1} v(s) ds - t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s) s^{\alpha-1} v(s) ds \right).$$

To show that  $M$  is compact on  $\mathcal{B}$  it is sufficient to show that  $\{(\hat{M}(v))' : v \in \hat{K}\}$  is uniformly bounded and equicontinuous on  $[0, 1]$ . We provide the details for equicontinuity as the details for uniform boundedness are straightforward.

Assume  $|p|_0 = P$  and assume  $|v|_0 \leq L$ . Let  $\epsilon > 0$ . As in the calculations above for  $g'$ ,  $(\hat{M}(v))'(0) = 0$  and

$$|(\hat{M}(v))'(t)| \leq \left( \frac{\alpha-1}{\alpha} + \frac{1}{\alpha-1} \right) PLt^{\alpha-1}.$$

Thus, there exists  $\delta_1 > 0$  such that if  $|t| < \delta_1$  then  $|(\hat{M}(v))'(t)| < \frac{\epsilon}{2}$ .

On  $[\delta_1, 1]$ ,  $\{(\hat{M}(v))' : v \in \hat{K}\}$  is shown to be equicontinuous by showing that  $\{(\hat{M}(v))'' : v \in \hat{K}\}$  is uniformly bounded. Now

$$\begin{aligned} (\hat{M}(v))'(t) &= (1-\alpha)t^{-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s) u(s) ds \\ &\quad + (\alpha-1)t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha)} p(s) u(s) ds, \end{aligned}$$

and so

$$\begin{aligned} (\hat{M}(v))''(t) &= -\alpha(1-\alpha)t^{-\alpha-1} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s) s^{\alpha-1} v(s) ds \\ &\quad - (\alpha-1)^2 t^{-\alpha} \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha)} p(s) s^{\alpha-1} v(s) ds \\ &\quad - (\alpha-1)^2 t^{-\alpha} \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha)} p(s) s^{\alpha-1} v(s) ds \\ &\quad + (\alpha-1)(\alpha-2)t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-3}}{\Gamma(\alpha-1)} p(s) s^{\alpha-1} v(s) ds. \end{aligned}$$

Each of the four terms can be bounded by a constant multiple of  $t^{\alpha-2}$ .

For the first term, notice

$$\left| t^{-\alpha-1} \int_0^t (t-s)^{\alpha-1} p(s) s^{\alpha-1} v(s) ds \right| \leq PLt^{-\alpha-1} \left| \int_0^t (t-s)^{\alpha-1} s^{\alpha-1} ds \right|.$$

Set  $s = rt$ . So

$$\begin{aligned} t^{-\alpha-1} \left| \int_0^t (t-s)^{\alpha-1} s^{\alpha-1} ds \right| &= t^{-\alpha-1} t^{\alpha-1} t^{\alpha-1} t \left| \int_0^1 (1-r)^{\alpha-1} r^{\alpha-1} dr \right| \\ &= t^{\alpha-2} |B(\alpha, \alpha)|, \end{aligned}$$

where  $B$  denotes the beta function.

In dealing with the second and third terms, first note

$$\left| t^{-\alpha} \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha)} p(s) s^{\alpha-1} v(s) ds \right| \leq PLt^{-\alpha} \left| \int_0^t (t-s)^{\alpha-2} s^{\alpha-1} ds \right|.$$

Set  $s = rt$ . Then

$$\begin{aligned} t^{-\alpha} \left| \int_0^t (t-s)^{\alpha-2} s^{\alpha-1} ds \right| &= t^{-\alpha} t^{\alpha-2} t^{\alpha-1} t \left| \int_0^1 (1-r)^{\alpha-2} r^{\alpha-1} dr \right| \\ &= t^{\alpha-2} |B(\alpha, \alpha-1)|. \end{aligned}$$

Notice  $B(\alpha, \alpha-1)$  is well-defined since  $1 < \alpha \leq 2$ .

Last, we obtain an analogous estimate for the fourth term. If  $\alpha = 2$ , this term is zero. If  $1 < \alpha < 2$ , first integrate by parts to obtain

$$\int_0^t (t-s)^{\alpha-3} s^{\alpha-1} ds = \frac{\alpha-1}{\alpha-2} \int_0^t (t-s)^{\alpha-2} s^{\alpha-2} ds.$$

Thus,

$$\begin{aligned} \left| t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-3}}{\Gamma(\alpha-1)} p(s) s^{\alpha-1} v(s) ds \right| &\leq PL t^{1-\alpha} \left| \int_0^t (t-s)^{\alpha-3} s^{\alpha-1} ds \right| \\ &= PL t^{1-\alpha} \left| \frac{\alpha-1}{\alpha-2} \int_0^t (t-s)^{\alpha-2} s^{\alpha-2} ds \right|. \end{aligned}$$

Again, by setting  $s = rt$ , we obtain

$$\begin{aligned} t^{1-\alpha} \left| \int_0^t (t-s)^{\alpha-2} s^{\alpha-2} ds \right| &= t^{1-\alpha} t^{\alpha-2} t^{\alpha-2} t \left| \int_0^1 (1-r)^{\alpha-2} r^{\alpha-2} dr \right| \\ &= t^{\alpha-2} |B(\alpha-1, \alpha-1)|. \end{aligned}$$

Again,  $B(\alpha-1, \alpha-1)$  is well-defined since  $1 < \alpha < 2$ . Therefore, if  $\alpha \neq 2$ ,

$$\begin{aligned} |(\hat{M}(v))''(t)| &\leq PL \left[ \frac{\alpha(\alpha-1)|B(\alpha, \alpha)|}{\Gamma(\alpha)} + \frac{2(\alpha-1)^2|B(\alpha, \alpha-1)|}{\Gamma(\alpha)} \right. \\ &\quad \left. + \frac{(\alpha-1)^2|B(\alpha-1, \alpha-1)|}{\Gamma(\alpha-1)} \right] t^{\alpha-2}, \end{aligned}$$

and if  $\alpha = 2$ ,

$$|(\hat{M}(v))''(t)| \leq \frac{4PL}{3}.$$

So  $\{(\hat{M}(v))'' : v \in \hat{K}\}$  is uniformly bounded on  $[\delta_1, 1]$ .

Since  $\{(\hat{M}(v))'' : v \in \hat{K}\}$  is uniformly bounded on  $[\delta_1, 1]$ , there exists  $\delta_2 > 0$  such that if  $|t_1 - t_2| < \delta_2$ ,  $t_1, t_2 \in [\delta_1, 1]$ , then  $|(\hat{M}(v))'(t_1) - (\hat{M}(v))'(t_2)| < \frac{\epsilon}{2}$ .

Set  $\delta = \min\{\delta_1, \delta_2\}$ . If  $|t_1 - t_2| < \delta$ ,  $t_1, t_2 \in [0, \delta_1]$ , then

$$|(\hat{M}(v))'(t_1) - (\hat{M}(v))'(t_2)| \leq |(\hat{M}(v))'(t_1)| + |(\hat{M}(v))'(t_2)| < \epsilon.$$

If  $|t_1 - t_2| < \delta$ ,  $t_1, t_2 \in [\delta_1, 1]$ , then

$$|(\hat{M}(v))'(t_1) - (\hat{M}(v))'(t_2)| \leq \frac{\epsilon}{2} < \epsilon.$$

If  $|t_1 - t_2| < \delta$ ,  $0 \leq t_1 < \delta_1 \leq t_2 \leq 1$ , then

$$\begin{aligned} &|(\hat{M}(v))'(t_1) - (\hat{M}(v))'(t_2)| \\ &\leq |(\hat{M}(v))'(t_1) - (\hat{M}(v))'(\delta_1)| + |(\hat{M}(v))'(\delta_1) - (\hat{M}(v))'(t_2)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Details for the operator  $N$  are similar and the proof is complete.  $\square$

Define the cone

$$\mathcal{P} = \{u \in \mathcal{B} : u(t) \geq 0 \text{ for } t \in [0, 1]\}.$$

**Lemma 3.2.** *The cone  $\mathcal{P}$  is solid in  $B$  and hence reproducing.*

*Proof.* Define

$$\Omega := \{u \in \mathcal{B} \mid u(t) > 0 \text{ for } t \in (0, 1), v(0) > 0, v'(1) < 0, \text{ where } u = t^{\alpha-1}v\}. \quad (3.4)$$

We will show  $\Omega \subset \mathcal{P}^\circ$ . Let  $u \in \Omega$ . Since  $v(0) > 0$ , there exists an  $\epsilon_1 > 0$  such that  $v(0) - \epsilon_1 > 0$ . Since  $v \in C^1[0, 1]$ , there exists an  $a \in (0, 1)$  such that  $v(t) > \epsilon_1$  for all  $t \in (0, a)$ . So  $u(t) = t^{\alpha-1}v(t) > \epsilon_1 t^{\alpha-1}$  for all  $t \in (0, a)$ . Now, since  $v'(1) < 0$ , there exists an  $\epsilon_2 > 0$  such that  $v'(1) + \epsilon_2 < 0$ . Then, since  $v(1) = 0$  and  $-v'(1) > \epsilon_2$ , there exists a  $b \in (a, 1)$  such that  $v(t) > (1-t)\epsilon_2$  for all  $t \in (b, 1]$ . Thus  $u(t) > b^{\alpha-1}(1-t)\epsilon_2$  for all  $t \in (b, 1]$ . Also, since  $u(t) > 0$  on  $[a, b]$ , there exists an  $\epsilon_3 > 0$  such that  $u(t) - \epsilon_3 > 0$  for all  $t \in [a, b]$ .

Let  $\epsilon = \min\{\frac{\epsilon_1}{2}, \frac{b^{\alpha-1}\epsilon_2}{2}, \frac{\epsilon_3}{2}\}$ . Define  $B_\epsilon(u) = \{\hat{u} \in \mathcal{B} : \|u - \hat{u}\| < \epsilon\}$ . Let  $\hat{u} \in B_\epsilon(u)$ . So  $\hat{u} = t^{\alpha-1}\hat{v}$ , where  $\hat{v} \in C^1[0, 1]$  with  $\hat{v}(1) = 0$ . Now

$$|\hat{u}(t) - u(t)| \leq t^{\alpha-1}\|\hat{u} - u\| < \epsilon t^{\alpha-1}.$$

So for  $t \in (0, a)$ ,  $\hat{u}(t) > u(t) - t^{\alpha-1}\epsilon > t^{\alpha-1}\epsilon_1 - t^{\alpha-1}\epsilon_1/2 = t^{\alpha-1}\epsilon_1/2$ . So  $\hat{u}(t) > 0$  for  $t \in (0, a)$ . By the Mean Value Theorem, for  $t \in (b, 1)$ ,  $|\hat{u}(t) - u(t)| \leq (1-t)\|\hat{u} - u\| < (1-t)\epsilon$ . So for  $t \in (b, 1)$ ,

$$\hat{u}(t) > u(t) - (1-t)\epsilon > b^{\alpha-1}(1-t)\epsilon_2 - (1-t)b^{\alpha-1}\epsilon_2/2 = (1-t)b^{\alpha-1}\epsilon_2/2.$$

So for  $t \in (b, 1)$ ,  $\hat{u}(t) > 0$ . Also,  $|\hat{u}(t) - u(t)| \leq \|\hat{u} - u\| < \epsilon$ . So for  $t \in [a, b]$ ,  $\hat{u}(t) > u(t) - \epsilon > \epsilon_3 - \epsilon_3/2 > 0$ . So  $\hat{u}(t) > 0$  for all  $t \in [a, b]$ . So  $\hat{u} \in \mathcal{P}$  and thus  $B_\epsilon(u) \subset \mathcal{P}$ . So  $\Omega \subset \mathcal{P}^\circ$ .  $\square$

**Lemma 3.3.** *The bounded linear operators  $M$  and  $N$  are  $u_0$ -positive with respect to  $\mathcal{P}$ .*

*Proof.* First, we show  $M : \mathcal{P} \setminus \{0\} \rightarrow \Omega \subset \mathcal{P}^\circ$ . Let  $u \in \mathcal{P}$ . So  $u(t) \geq 0$ . Then since  $G(t, s) \geq 0$  on  $[0, 1] \times [0, 1]$  and  $p(t) \geq 0$  on  $[0, 1]$ ,

$$Mu(x) = \int_0^1 G(t, s)p(s)u(s)ds \geq 0,$$

for  $0 \leq t \leq 1$ . So  $M : \mathcal{P} \rightarrow \mathcal{P}$ .

Now let  $u \in \mathcal{P} \setminus \{0\}$ . So there exists a compact interval  $[\alpha, \beta] \subset [0, 1]$  such that  $u(t) > 0$  and  $p(t) > 0$  for all  $t \in [\alpha, \beta]$ . Then, since  $G(t, s) > 0$  on  $(0, 1) \times (0, 1)$ ,

$$\begin{aligned} Mu(t) &= \int_0^1 G(t, s)p(s)u(s)ds \\ &\geq \int_\alpha^\beta G(t, s)p(s)u(s)ds > 0, \end{aligned}$$

for  $0 < t < 1$ . Now

$$Mu(t) = t^{\alpha-1} \left( \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds - t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds \right).$$

Let

$$v(t) = \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds - t^{1-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds.$$



So  $v(0) = \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds > 0$  and

$$v'(1) = -(1-\alpha) \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} p(s)u(s)ds - (\alpha-1) \int_0^1 \frac{(1-s)^{\alpha-2}}{\Gamma(\alpha)} p(s)u(s)ds < 0.$$

So  $M : \mathcal{P} \setminus \{0\} \rightarrow \Omega \subset \mathcal{P}^\circ$ .

Now choose any  $u_0 \in \mathcal{P} \setminus \{0\}$ , and let  $u \in \mathcal{P} \setminus \{0\}$ . So  $Mu \in \Omega \subset \mathcal{P}^\circ$ . Choose  $k_1 > 0$  sufficiently small and  $k_2$  sufficiently large so that  $Mu - k_1u_0 \in \mathcal{P}^\circ$  and  $u_0 - \frac{1}{k_2}Mu \in \mathcal{P}^\circ$ . So  $k_1u_0 \leq Mu$  with respect to  $\mathcal{P}$  and  $Mu \leq k_2u_0$  with respect to  $\mathcal{P}$ . Thus  $k_1u_0 \leq Mu \leq k_2u_0$  with respect to  $\mathcal{P}$  and so  $M$  is  $u_0$ -positive with respect to  $\mathcal{P}$ . Similarly,  $N$  is  $u_0$ -positive.  $\square$

**Remark 3.4.** Notice that

$$\Lambda u = Mu = \int_0^1 G(t,s)p(s)u(s)ds,$$

if and only if

$$u(t) = \frac{1}{\Lambda} \int_0^1 G(t,s)p(s)u(s)ds,$$

if and only if

$$D_{0+}^\alpha u(t) + \frac{1}{\Lambda} p(t)u(t) = 0, \quad 0 < t < 1,$$

with  $u(0) = u(1) = 0$ .

So the eigenvalues of (1.1),(1.3) are reciprocals of eigenvalues of  $M$ , and conversely. Similarly, eigenvalues of (1.2),(1.3) are reciprocals of eigenvalues of  $N$ , and conversely.

**Theorem 3.5.** *Let  $\mathcal{B}$ ,  $\mathcal{P}$ ,  $M$ , and  $N$  be defined as earlier. Then  $M$  (and  $N$ ) has an eigenvalue that is simple, positive, and larger than the absolute value of any other eigenvalue, with an essentially unique eigenvector that can be chosen to be in  $\mathcal{P}^\circ$ .*

*Proof.* Since  $M$  is a compact linear operator that is  $u_0$ -positive with respect to  $\mathcal{P}$ , by Theorem 2.7,  $M$  has an essentially unique eigenvector, say  $u \in \mathcal{P}$ , and eigenvalue  $\Lambda$  with the above properties. Since  $u \neq 0$ ,  $Mu \in \Omega \subset \mathcal{P}^\circ$  and  $u = M(\frac{1}{\Lambda}u) \in \mathcal{P}^\circ$ .  $\square$

**Theorem 3.6.** *Let  $\mathcal{B}$ ,  $\mathcal{P}$ ,  $M$ , and  $N$  be defined as earlier. Let  $p(t) \leq q(t)$  on  $[0, 1]$ . Let  $\Lambda_1$  and  $\Lambda_2$  be the eigenvalues defined in Theorem 3.5 associated with  $M$  and  $N$ , respectively, with the essentially unique eigenvectors  $u_1$  and  $u_2 \in \mathcal{P}^\circ$ . Then  $\Lambda_1 \leq \Lambda_2$ , and  $\Lambda_1 = \Lambda_2$  if and only if  $p(t) = q(t)$  on  $[0, 1]$ .*

*Proof.* Let  $p(t) \leq q(t)$  on  $[0, 1]$ . So for any  $u \in \mathcal{P}$  and  $t \in [0, 1]$ ,

$$(Nu - Mu)(t) = \int_0^1 G(t,s)(q(s) - p(s))u(s)ds \geq 0.$$

So  $Nu - Mu \in \mathcal{P}$  for all  $u \in \mathcal{P}$ , or  $M \leq N$  with respect to  $\mathcal{P}$ . Then by Theorem 2.8,  $\Lambda_1 \leq \Lambda_2$ .

If  $p(t) = q(t)$ , then  $\Lambda_1 = \Lambda_2$ . Now suppose  $p(t) \neq q(t)$ . So  $p(t) < q(t)$  on some subinterval  $[\alpha, \beta] \subset [0, 1]$ . Then  $(N - M)u_1 \in \Omega \subset \mathcal{P}^\circ$  and so there exists  $\epsilon > 0$  such that  $(N - M)u_1 - \epsilon u_1 \in \mathcal{P}$ . So  $\Lambda_1 u_1 + \epsilon u_1 = Mu_1 + \epsilon u_1 \leq Nu_1$ , implying  $Nu_1 \geq (\Lambda_1 + \epsilon)u_1$ . Since  $N \leq N$  and  $Nu_2 = \Lambda_2 u_2$ , by Theorem 2.8,  $\Lambda_1 + \epsilon \leq \Lambda_2$ , or  $\Lambda_1 < \Lambda_2$ .  $\square$

By Remark 3.4, the following theorem is an immediate consequence of Theorems 3.5 and 3.6.

**Theorem 3.7.** *Assume the hypotheses of Theorem 3.6. Then there exists smallest positive eigenvalues  $\lambda_1$  and  $\lambda_2$  of (1.1),(1.3) and (1.2),(1.3), respectively, each of which is simple, positive, and less than the absolute value of any other eigenvalue of the corresponding problems. Also, eigenfunctions corresponding to  $\lambda_1$  and  $\lambda_2$  may be chosen to belong to  $\mathcal{P}^\circ$ . Finally,  $\lambda_1 \geq \lambda_2$ , and  $\lambda_1 = \lambda_2$  if and only if  $p(t) = q(t)$  for all  $t \in [0, 1]$ .*

#### REFERENCES

- [1] R. P. Agarwal, M. Benchohra, S. Hamani; *A survey on existence results for boundary value problems of nonlinear fractional differential equations and inclusions*. Acta. Appl. Math. **109** (2010), 973-1003.
- [2] M. Al-Refai; *Basic results on nonlinear eigenvalue problems of fractional order*. Electron. J. Differential Equations **2012** (2012), No. 191, 1-12.
- [3] Z. Bia, H. Lu; *Positive solutions for boundary value problem of nonlinear fractional differential equation*. J. Math. Anal. Appl. **311** (2005), 495-505.
- [4] C. J. Chyan, J. M. Davis, J. Henderson, W. K. C. Yin; *Eigenvalue comparisons for differential equations on a measure chain*. Electron. J. Differential Eqns. **1998** (1998), No. 35, 1-7.
- [5] J. M. Davis, P. W. Eloe, J. Henderson; *Comparison of eigenvalues for discrete Lidstone boundary value problems*. Dyn. Sys. Appl. **8** (1999), 381-388.
- [6] K. Diethelm; *The Analysis of Fractional Differential equations. An Application-oriented Exposition Using Differential Operators of Caputo Type*. Lecture Notes in Mathematics, 2004, Springer-Verlag, Berlin, 2010.
- [7] P. W. Eloe, J. Henderson; *Comparison of eigenvalues for a class of two-point boundary value problems*. Appl. Anal. **34** (1989), 25-34.
- [8] P. W. Eloe, J. Henderson; *Comparison of eigenvalues for a class of multipoint boundary value problems*. Recent Trends in Ordinary Differential Equations **1** (1992), 179-188.
- [9] R. D. Gentry, C. C. Travis; *Comparison of eigenvalues associated with linear differential equations of arbitrary order*. Trans. Amer. Math. Soc. **223** (1967), 167-179.
- [10] X. Han, H. Gao; *Existence of positive solutions for eigenvalue problem of nonlinear fractional differential equations*. Adv. Difference Equ. **2012**, (2012), No. 66, 8 pp.
- [11] D. Hankerson, A. Peterson; *Comparison of eigenvalues for focal point problems for  $n$ th order difference equations*. Differential Integral Eqns. **3** (1990), 363-380.
- [12] J. Hoffacker; *Green's functions and eigenvalue comparisons for a focal problem on time scales*. Comput. Math. Appl., **45** (2003), 1339-1368.
- [13] M. Keener, C. C. Travis; *Positive cones and focal points for a class of  $n$ th order differential equations*. Trans. Amer. Math. Soc. **237** (1978), 331-351.
- [14] M. G. Krein and M. A. Rutman; *Linear operators leaving a cone invariant in a Banach space*. Translations Amer. Math. Soc., Series 1, Volume 10, 199-325, American Mathematical Society, Providence, RI, 1962.
- [15] M. Krasnosel'skii; *Positive Solutions of Operator Equations*. Fizmatgiz, Moscow, 1962; English Translation P. Noordhoff Ltd. Gronigen, The Netherlands, 1964.
- [16] J. T. Neugebauer; *Methods of extending lower order problems to higher order problems in the context of smallest eigenvalue comparisons*. Electron. J. Qual. Theory Differ. Equ. **99** (2011), 1-16.
- [17] S. Samko, A. Kilbas, O. Marichev; *Fractional Integrals and Derivatives: Theory and Applications*. Gordon and Breach, Yverdon, 1993.
- [18] C. C. Travis; *Comparison of eigenvalues for linear differential equations*. Proc. Amer. Math. Soc. **96** (1986), 437-442.
- [19] J. Wu, X. Zhang; *Eigenvalue problem of nonlinear semipositone higher order fractional differential equations*. Abst. Appl. Anal. **2012** (2012), Art. ID 740760, 14 pp.
- [20] X. Zhang, L. Liu, B. Wiwatanapataphee, Y. Wu; *Positive solutions of eigenvalue problems for a class of fractional differential equations with derivatives*. Abst. Appl. Anal. **2012** (2012), Art. ID 512127, 16 pp.

PAUL W. ELOE

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DAYTON, DAYTON, OH 45469, USA

*E-mail address:* `peloe1@udayton.edu`

JEFFREY T. NEUGEBAUER

DEPARTMENT OF MATHEMATICS AND STATISTICS, EASTERN KENTUCKY UNIVERSITY, RICHMOND,  
KY 40475, USA

*E-mail address:* `jeffrey.neugebauer@eku.edu`