

2011

Optimal intervals for uniqueness of solutions for nonlocal boundary value problems

Paul W. Eloë

University of Dayton, peloe1@udayton.edu

Johnny Henderson

Baylor University

Follow this and additional works at: https://ecommons.udayton.edu/mth_fac_pub



Part of the [Mathematics Commons](#)

eCommons Citation

Eloë, Paul W. and Henderson, Johnny, "Optimal intervals for uniqueness of solutions for nonlocal boundary value problems" (2011). *Mathematics Faculty Publications*. 127.

https://ecommons.udayton.edu/mth_fac_pub/127

This Article is brought to you for free and open access by the Department of Mathematics at eCommons. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of eCommons. For more information, please contact frice1@udayton.edu, mschlangen1@udayton.edu.

INTERNATIONAL PUBLICATIONS (USA)

Communications on Applied Nonlinear Analysis
Volume 18(2011), Number 3, 89–97

Optimal Intervals for Uniqueness of Solutions for Nonlocal Boundary Value Problems

Paul W. Eloë
University of Dayton
Department of Mathematics
Dayton, Ohio 45469-2316 USA
Paul.Eloe@notes.udayton.edu

Johnny Henderson
Baylor University
Department of Mathematics
Waco, Texas 76798-7328 USA
Johnny_Henderson@baylor.edu

(In Memory of Professor Louis J. Grimm, November 30, 1933 – December 14, 2010)

Communicated by the Editors
(Received June 3, 2011; Accepted July 14, 2011)

Abstract

For the n th order differential equation, $y^{(n)} = f(t, y, y', \dots, y^{(n-1)})$, where $f(t, r_1, r_2, \dots, r_n)$ is Lipschitz continuous in terms of $r_i, 1 \leq i \leq n$, we obtain optimal bounds on the length of intervals on which solutions are unique for certain nonlocal three point boundary value problems. These bounds are obtained through an application of the Pontryagin Maximum Principle from the theory of optimal control.

Key words: Nonlocal boundary value problem, optimal length intervals, Pontryagin Maximum Principle.

AMS Subject Classification: 34B15

1 Introduction

In this paper, we shall be concerned with the n th order differential equation,

$$y^{(n)} = f(t, y, y', \dots, y^{(n-1)}), \quad a < t < b, \quad (1.1)$$

for which the assumptions in the following hypothesis hold throughout.

Hypothesis. $f(t, r_1, \dots, r_n) : (a, b) \times \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous, and for nonnegative constants $k_i, 1 \leq i \leq n$, satisfies the Lipschitz condition,

$$|f(t, r_1, \dots, r_n) - f(t, s_1, \dots, s_n)| \leq \sum_{i=1}^n k_i |r_i - s_i|, \quad (1.2)$$

for each $(t, r_1, \dots, r_n), (t, s_1, \dots, s_n) \in (a, b) \times \mathbb{R}^n$.

We characterize optimal length for subintervals of (a, b) , in terms of the Lipschitz coefficients $k_i, 1 \leq i \leq n$, on which solutions are unique for problems involving (1.1) and satisfying the *nonlocal* three point boundary conditions,

$$y^{(i-1)}(t_1) = y_i, \quad 1 \leq i \leq n-1, \quad y^{(n-2)}(t_2) - y^{(n-2)}(t_3) = y_n, \quad (1.3)$$

where $a < t_1 < t_2 < t_3 < b$, and $y_1, \dots, y_n \in \mathbb{R}$.

More precisely, we characterize optimal length for subintervals of (a, b) on which solutions of (1.1), (1.3) are unique. Such uniqueness results are of interest, because in many cases, uniqueness of solutions implies existence of solutions for boundary value problems; see, for example, the papers [5, 7, 9, 18, 19, 22, 24, 32] and the references therein.

There is a close connection between the boundary value problem (1.1), (1.3) and certain right focal boundary value problems for (1.1). Because of this relationship, we will eventually establish that it suffices for us to characterize optimal length subintervals of (a, b) on which solutions are unique for (1.1) satisfying the right focal boundary conditions,

$$y^{(i-1)}(t_1) = y_i, \quad 1 \leq i \leq n-1, \quad y^{(n-1)}(t_2) = y_n, \quad (1.4)$$

where $a < t_1 < t_2 < b$, and $y_1, \dots, y_n \in \mathbb{R}$. The connection between this characterization and the characterization for our three point nonlocal problems is through a simple application of the Mean Value Theorem.

Theorem 1.1 *If solutions for (1.1), (1.4) are unique, when they exist on (a, b) , then solutions for (1.1), (1.3) are unique, when they exist on (a, b) .*

Thus, in view of Theorem 1.1, conditions sufficient to provide uniqueness of solutions, when they exist on (a, b) , for two point right focal boundary value problems (1.1), (1.4), are sufficient to provide uniqueness of solutions, when they exist on (a, b) for three point nonlocal boundary value problems (1.1), (1.3).

Our process will involve development of a scenario in which the Pontryagin Maximum Principle can be applied. We follow a pattern that has an extensive history, with first motivation found in the papers by Melentsova [36] and Melentsova and Mil'shtein [37, 38].

Those papers were subsequently adapted to the context of several types of boundary value problems by Jackson [28, 29], Eloe and Henderson [8], Hankerson and Henderson [17] and Henderson *et al.* [6, 20, 21, 23, 25].

Interest in nonlocal boundary value problems also has a long history, both in application and theory, as can be seen in this list of papers and the references therein: [1] -[4], [10, 11], [13] - [16], [23], [26, 27], [30, 31], [34, 35], [39] - [47].

2 Optimal Intervals for Uniqueness of Solutions

In this section, we apply the Pontryagin Maximum Principle to obtain a characterization, in terms of the Lipschitz constants k_i , $1 \leq i \leq n$, for the optimal length of subintervals of (a, b) on which solutions are unique, when they exist for the right focal boundary value problem (1.1), (1.4). This length, it will be argued later, is optimal for uniqueness of solutions for the three point nonlocal boundary value problem (1.1), (1.3).

We first introduce a set of vector-valued *control functions*

$$\mathcal{U} := \{ \mathbf{v}(t) = (v_1(t), \dots, v_n(t))^T \in \mathbb{R}^n \mid v_i(t) \text{ are Lebesgue measurable and } |v_i(t)| \leq k_i \text{ on } (a, b), i = 1, \dots, n \}.$$

We will be concerned with boundary value problems associated with linear differential equations of the form

$$x^{(n)} = \sum_{i=1}^n u_i(t)x^{(i-1)}, \quad (2.1)$$

where $\mathbf{u}(t) = (u_1(t), \dots, u_n(t))^T \in \mathcal{U}$. We immediately make a connection of these linear differential equations with solutions of (1.1), (1.4). Much of our analysis will be based upon our choosing, if they exist, distinct solutions $y(t)$ and $z(t)$ of (1.1), (1.4).

If $y(t)$ and $z(t)$ are distinct solutions of (1.1), (1.4), then their difference $x(t) := y(t) - z(t)$ satisfies

$$x^{(i-1)}(t_1) = x^{(n-1)}(t_2) = 0, \quad 1 \leq i \leq n-1, \quad (2.2)$$

for some $a < t_1 < t_2 < b$, and if for $1 \leq i \leq n$,

$$u_i(t) := \begin{cases} \frac{f(t, z(t), \dots, z^{(i-2)}(t), y^{(i-1)}(t), \dots, y^{(n-1)}(t)) - f(t, z(t), \dots, z^{(i-1)}(t), y^{(i)}(t), \dots, y^{(n-1)}(t))}{y^{(i-1)}(t) - z^{(i-1)}(t)}, & y^{(i-1)}(t) \neq z^{(i-1)}(t), \\ 0, & y^{(i-1)}(t) = z^{(i-1)}(t), \end{cases}$$

then $u_i(t)$ is Lebesgue measurable, and $|u_i(t)| \leq k_i$, $i = 1, \dots, n$, so that $\mathbf{u}(t) = (u_1(t), \dots, u_n(t))^T \in \mathcal{U}$, and $x(t)$ is a nontrivial solution of the boundary value problem (2.1), (2.2). It follows from optimal control theory (cf. Gamkrelidze [12, p. 147] and Lee and Markus [33, p. 259]), there is a boundary value problem in the class (2.1), (2.2), which has a nontrivial time optimal solution; that is, there exists at least one nontrivial $\mathbf{u}^* \in \mathcal{U}$ and points $t_1 \leq c < d \leq t_2$ such that

$$x^{(n)} = \sum_{i=1}^n u_i^*(t)x^{(i-1)}, \quad (2.3)$$

$$x^{(i-1)}(c) = x^{(n-1)}(d) = 0, \quad 1 \leq i \leq n, \quad (2.4)$$

has a nontrivial solution, $x_0(t)$, and $d - c$ is a minimum over all such solutions. For this time optimal solution, $x_0(t)$, set $\mathbf{x}_0(t) = (x_0(t), \dots, x_0^{(n-1)}(t))^T$. Then $\mathbf{x}_0(t)$ is a solution of a first order system,

$$\mathbf{r}' = A[\mathbf{u}^*(t)]\mathbf{r}.$$

By the Pontryagin Maximum Principle, the adjoint system, whose form is given by

$$\mathbf{x}' = -A^T[\mathbf{u}^*(t)]\mathbf{x}, \quad a < t < b, \quad (2.5)$$

has a nontrivial solution, $\mathbf{x}^*(t) = (x_1^*(t), \dots, x_n^*(t))^T$ such that, for *a. e.* $t \in [c, d]$,

- (i) $\sum_{i=1}^n x_0^{(i)}(t)x_i^*(t) = \langle \mathbf{x}'_0(t), \mathbf{x}^*(t) \rangle = \max_{\mathbf{u} \in \mathcal{U}} \{ \langle A[\mathbf{u}(t)]\mathbf{x}_0(t), \mathbf{x}^*(t) \rangle \}$,
- (ii) $\langle \mathbf{x}'_0(t), \mathbf{x}^*(t) \rangle$ is a nonnegative constant,
- (iii) $x_n^*(c) = x_1^*(d) = \dots = x_{n-1}^*(d) = 0$.

The maximum condition in (i) can be rewritten as

$$x_n^*(t) \sum_{i=1}^n u_i^*(t) x_0^{(i-1)}(t) = \max_{\mathbf{u} \in \mathcal{U}} \left\{ x_n^*(t) \sum_{i=1}^n u_i(t) x_0^{(i-1)}(t) \right\}, \quad (2.6)$$

for *a. e.* $t \in [c, d]$.

By its time optimality and repeated applications of Rolle's Theorem, $x_0(t) \neq 0$, $t \in (c, d]$. In fact, for each $1 \leq i \leq n$, $x_0^{(i-1)}(t) \neq 0$ on (c, d) . We may assume without loss of generality that $x_0(t) > 0$ on $(c, d]$. If $x_n^*(t)$ has no zeros on (c, d) , then we can use (2.6) to determine an optimal control $\mathbf{u}^*(t)$, for *a. e.* $t \in [c, d]$. We now examine the sign of $x_n^*(t)$ on (c, d) .

In that direction, if $\bar{\mathbf{u}} \in \mathcal{U}$ is such that the boundary value problem given by (2.1) and (2.2), for some $a < t_1 < t_2 < b$, has a nontrivial solution, then the adjoint system

$$\alpha' = -A^T[\bar{\mathbf{u}}(t)]\alpha, \quad t \in (a, b), \quad (2.7)$$

$$\alpha_n(t_1) = \alpha_1(t_2) = \dots = \alpha_{n-1}(t_2) = 0, \quad (2.8)$$

also has a nontrivial solution, and conversely. That is, the Pontryagin Maximum Principle associates with a time optimal solution of boundary value problem (2.1), (2.2) a time optimal solution of boundary value problem (2.7), (2.8), and conversely. Hence, it follows by its own time optimality that $x_n^*(t)$ does not vanish on (c, d) .

Now, $x_0(t) > 0$ on $(c, d]$, and so we have from (2.6) that, if $x_n^*(t) < 0$ on (c, d) , then the time optimal solution $x_0(t)$ is a solution of

$$x^{(n)} = -k_1 x - \sum_{i=2}^n k_i |x^{(i-1)}| \quad (2.9)$$

on $[c, d]$, while if $x_n^*(t) > 0$ on (c, d) , then the time optimal solution $x_0(t)$ is a solution of

$$x^{(n)} = k_1 x + \sum_{i=2}^n k_i |x^{(i-1)}| \quad (2.10)$$

on $[c, d]$. Since $n - (n - 1) = 1$ is odd, the result by Jackson [29, Theorem 2] yields that $x_n^*(t) < 0$ on (c, d) , so that $x_0(t)$ is a solution of (2.9) on $[c, d]$. Moreover, from the assumed positivity of $x_0(t)$ and the nature of the boundary conditions (2.4), along with the fact that $x_0^{(i-1)}(t) \neq 0$ on (c, d) , $1 \leq i \leq n$, it follows that $x_0^{(i-1)}(t) > 0$ on (c, d) , $1 \leq i \leq n$. As a consequence, not only is $x_0(t)$ a solution of (2.9), but also where (2.9) takes the form

$$x^{(n)} = - \sum_{i=1}^n k_i x^{(i-1)}. \quad (2.11)$$

Recall that our discussion is based on (1.1) having distinct solutions whose difference satisfies (2.2). In addition, if sufficient sign conditions are satisfied by the optimal solution $x_0(t)$ of the boundary value problem (2.1), (2.2) and by the component $x_n^*(t)$ of the solution of the associated adjoint system (2.5), then optimal intervals can be determined on which only trivial solutions exist for boundary value problems (2.9), (2.2) or (2.10), (2.2). Ultimately, a more detailed sign analysis led to determination of optimal intervals on which only trivial solutions exist for only the boundary value problem (2.11), (2.2). As a consequence, solutions of the boundary value problem (1.1), (1.4) will be unique on such subintervals.

Theorem 2.1 *If there is a vector-valued $\mathbf{u}(t) \in \mathcal{U}$ for all $a < t < b$, for which the boundary value problem (2.1), (2.2) has a nontrivial solution for some $a < t_1 < t_2 < b$, and if $x_0(t)$ is a time optimal solution satisfying (2.4), where $d - c$ is a minimum, then $x_0(t)$ is a solution of (2.11) on $[c, d]$.*

Theorem 2.2 *Let $\ell = \ell(k_1, \dots, k_n) > 0$ be the smallest positive number such that there exists a solution $x(t)$ of the boundary value problem for (2.11) satisfying*

$$x^{(i-1)}(0) = 0, \quad 1 \leq i \leq n-1, \quad x^{(n-1)}(\ell) = 0, \quad (2.12)$$

with $x(t) > 0$ on $(0, \ell]$, or $\ell = \infty$ if no such solution exists. If $y(t)$ and $z(t)$ are solutions of the boundary value problem (1.1), (1.4), for some $a < t_1 < t_2 < b$, and if $t_2 - t_1 < \ell$, it follows that $y(t) \equiv z(t)$ on $[t_1, t_2]$, and this is best possible for the class of all differential equations satisfying the Lipschitz condition (1.2).

Proof: Since equation (2.11) is autonomous, translations of solutions are again solutions of (2.11). Hence, it suffices to apply Theorem 2.1 with respect to the boundary conditions at 0 and ℓ .

Now, if $y(t)$ and $z(t)$ are distinct solutions of (1.1) whose difference $w(t) := y(t) - z(t)$ satisfies (2.2), where $t_2 - t_1 < \ell$, then $w(t)$ is a nontrivial solution of the boundary value problem (2.1), (2.2), for appropriately defined $\mathbf{u} \in \mathcal{U}$. Then, from the discussion and Theorem 2.1, equation (2.11) has a nontrivial solution on a subinterval of length less than ℓ . But, by the minimality of ℓ , such a boundary value problem can have only the trivial solution; this is a contradiction. Therefore, solutions of the boundary value problem (1.1), (1.4) are unique, whenever $t_2 - t_1 < \ell$.

That this is best possible from the fact that (2.11) satisfies the Lipschitz condition (1.2), and if $\ell \neq \infty$, then $x(t)$ is a nontrivial solution of (2.11) and (2.2) on $[0, \ell]$. The boundary value problem also has the trivial solution. \square

Remark 2.1 Since (2.11) is a linear equation, we observe that, if $x(t)$ is the solution, of the initial value problem for (2.11), satisfying,

$$x^{(i-1)}(0) = 0, \quad 1 \leq i \leq n-1, \quad x^{(n-1)}(0) = 1,$$

and if $\eta > 0$ is the first positive number such that $x^{(n-1)}(\eta) = 0$, then $\eta = \ell(k_1, \dots, k_n)$ of Theorem 2.2.

Because of the uniqueness relationships stated in Theorem 1.1, we can apply Theorem 2.2 to obtain optimal intervals for uniqueness of solutions of the boundary value problem (1.1), (1.3).

Theorem 2.3 Let ℓ be as in Theorem 2.2. If $y(t)$ and $z(t)$ are solutions of the boundary value problem (1.1), (1.3), for some $a < t_1 < t_2 < t_3 < b$, and if $t_3 - t_1 \leq \ell$, it follows that $y(t) \equiv z(t)$ on $[t_1, t_3]$, and this is best possible for the class of all differential equations satisfying the Lipschitz condition (1.2).

Proof: In view of Theorem 1.1 and Theorem 2.2, solutions of the boundary value problem (1.1), (1.3) are unique, when $t_3 - t_1 \leq \ell$. To see again that this is best possible, consider the nontrivial solution $x(t)$ of (2.11) and (2.12) in Theorem 2.2.

Let $\epsilon > 0$ be sufficiently small that $x(t)$ is a solution of (2.11) on $[0, \ell + \epsilon]$. Now, $x^{(n)}(t) < 0$ on $[0, \ell + \epsilon]$. From (2.12), $x^{(n-1)}(\ell) = 0$, and since $x^{(n)}(\ell) < 0$, we have that $x^{(n-2)}(t)$ has a positive maximum at ℓ . So, there exist $0 < \tau_1 < \ell < \tau_2 < \ell + \epsilon$ such that $x(t)$ is a nontrivial solution of (2.11) satisfying $x^{(i-1)}(0) = 0, 1 \leq i \leq n-1$, and $x^{(n-2)}(\tau_1) - x^{(n-2)}(\tau_2) = 0$. This boundary value problem also has the trivial solution. Since $\epsilon > 0$ was arbitrary, the “best possible” statement follows for uniqueness of solutions of the boundary value problem (1.1), (1.3). \square

3 Optimal Intervals of Existence for Linear Equations

In the case of boundary value problem (1.1), (1.3), we do not have a “uniqueness implies existence” theorem to appeal to, since this is an open question for this type of boundary value problem. However, uniqueness does imply existence for linear differential equations, and so the following corollary can be stated.

Corollary 3.1 Let ℓ be as in Theorem 2.2. Assume $p_i(t), 1 \leq i \leq n$, and $q(t)$ are continuous on (a, b) and that $|p_i(t)| \leq k_i$ on $(a, b), 1 \leq i \leq n$. If $a < t_1 < t_2 < t_3 < b$ and $t_3 - t_1 < \ell$, then the boundary value problem,

$$y^{(n)} = \sum_{i=1}^n p_i(t)y^{(i-1)} + q(t),$$

$$y^{(i-1)}(t_1) = y_i, \quad 1 \leq i \leq n-1, \quad y^{(n-2)}(t_2) - y^{(n-2)}(t_3) = y_n,$$

has a solution for any assignment of values of $y_i \in \mathbb{R}, 1 \leq i \leq n$.

References

- [1] B. Ahmad and J. J. Nieto, Existence of solutions for nonlocal boundary value problems of higher-order nonlinear fractional differential equations, *Abstr. Appl. Anal.* **2009**, Art. ID 494720, 9 pp.
- [2] M. Bahaj, Remarks on the existence results for second-order differential inclusions with nonlocal conditions, *J. Dyn. Control Syst.* **15** (2009), no. 1, 2–43.
- [3] C. Bai and J. Fang, Existence of multiple positive solutions for m -point boundary value problems, *J. Math. Anal. Appl.* **281** (2003), 76–85.
- [4] Z. N. Benbouziane, A. Boucherif and S. M. Bouguima, Third order nonlocal multipoint boundary value problems, *Dynam. Systems Appl.* **13** (2004), 41–48.
- [5] C. J. Chyan and J. Henderson, Uniqueness implies existence for (n, p) boundary value problems, *Appl. Anal.* **73** (1999), no. 3-4, 543–556.
- [6] S. Clark and J. Henderson, Optimal interval lengths for nonlocal boundary value problems associated with third order Lipschitz equations, *J. Math. Anal. Appl.* **322** (2006), 468–476.
- [7] S. Clark and J. Henderson, Uniqueness implies existence and uniqueness criterion for nonlocal boundary value problems for third order differential equations, *Proc. Amer. Math. Soc.* **134** (2006), no. 11, 3363–3372.
- [8] P. W. Eloe and J. Henderson, Optimal intervals for third order Lipschitz equations, *Differential Integral Equations* **2** (1989), 397–404.
- [9] P. W. Eloe and J. Henderson, Uniqueness implies existence and uniqueness conditions for nonlocal boundary value problems for n th order differential equations, *J. Math. Anal. Appl.* **331** (2007), no. 1, 240–247
- [10] W. Feng and J. R. L. Webb, Solvability of a three-point nonlinear boundary value problem at resonance, *Nonlinear Anal.* **30** (1997), 3227–3238.
- [11] W. Feng and J. R. L. Webb, Solvability of an m -point nonlinear boundary value problem with nonlinear growth, *J. Math. Anal. Appl.* **212** (1997), 467–480.
- [12] R. Gamkrelidze, *Principles of Optimal Control*, Plenum, New York, 1978.
- [13] J. R. Graef, J. Henderson and B. Yang, Existence of positive solutions of a higher order nonlocal singular boundary value problem, *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.* **16** (2009), Differential Equations and Dynamical Systems, suppl. S1, 147–152.
- [14] J. R. Graef and J. R. L. Webb, Third order boundary value problems with nonlocal boundary conditions, *Nonlinear Anal.* **71** (2009), no. 5-6, 1542–1551.
- [15] Y. Guo, W. Shan and W. Ge, Positive solutions for second-order m -point boundary value problems, *J. Comput. Appl.* **151** (2003), 415–424.

- [16] C. P. Gupta, S. K. Ntouyas and P. Ch. Tsamatos, Solvability of an m -point boundary value problem for second order ordinary differential equations, *J. Math. Anal. Appl.* **189** (1995), 575–584.
- [17] D. Hankerson and J. Henderson, Optimality for boundary value problems for Lipschitz equations, *J. Differential Equations* **77** (1989), 392–404.
- [18] P. Hartman, On n -parameter families and interpolation problems for nonlinear ordinary differential equations, *Trans. Amer. Math. Soc.* **154** (1971), 201–226.
- [19] J. Henderson, Existence of solutions of right focal point boundary value problems for ordinary differential equations, *Nonlinear Anal.* **5** (1981), no. 9, 989–1002.
- [20] J. Henderson, Best interval lengths for boundary value problems for third order Lipschitz equations, *SIAM J. Math. Anal.* **18** (1987), 293–305.
- [21] J. Henderson, Boundary value problems for n th order Lipschitz equations, *J. Math. Anal. Appl.* **144** (1988), 196–210.
- [22] J. Henderson, Uniqueness implies existence for three-point boundary value problems for second order differential equations, *Appl. Math. Lett.* **18** (2005), 905–909.
- [23] J. Henderson, Optimal interval lengths for nonlocal boundary value problems for second order Lipschitz equations, *Comm. Appl. Anal.*, in press.
- [24] J. Henderson, B. Karna and C. C. Tisdell, Existence of solutions for three-point boundary value problems for second order equations, *Proc. Amer. Math. Soc.* **133** (2005), 1365–1369.
- [25] J. Henderson and R. McGwier, Uniqueness, existence and optimality for fourth order Lipschitz equations, *J. Differential Equations* **67** (1987), 414–440.
- [26] E. Hernández, Existence of solutions for an abstract second-order differential equation with nonlocal conditions, *Electron. J. Differential Equations* **2009**, No. 96, 10 pp.
- [27] G. Infante, Nonlocal boundary value problems with two nonlinear boundary conditions, *Commun. Appl. Anal.* **12** (2008), no. 3, 279–288.
- [28] L. K. Jackson, Existence and uniqueness of solutions for boundary value problems for Lipschitz equations, *J. Differential Equations* **32** (1979), 76–90.
- [29] L. K. Jackson, Boundary value problems for Lipschitz equations, *Differential Equations (Proc. Eighth Fall Conf., Oklahoma State Univ., Stillwater, Okla., 1979)*, pp. 31–50, Academic Press, New York, 1980.
- [30] P. Kang and Z. Wei, Three positive solutions of singular nonlocal boundary value problems for systems of nonlinear second-order ordinary differential equations, *Nonlinear Anal.* **70** (2009), no. 1, 444–451.
- [31] R. A. Khan, Quasilinearization method and nonlocal singular three point boundary value problems, *Electron. J. Qual. Theory Differ. Equ.* **2009**, Special Edition I, No. 17, 13 pp.

- [32] G. Klaasen, Existence theorems for boundary value problems for n th order ordinary differential equations, *Rocky Mtn. J. Math.* **3** (1973), 457–472.
- [33] E. Lee and L. Markus, *Foundations of Optimal Control*, Wiley, New York, 1967.
- [34] M. Li and C. Kou, Existence results for second-order impulsive neutral functional differential equations with nonlocal conditions, *Discrete Dyn. Nat. Soc.* **2009**, Art. ID 641368, 11 pp.
- [35] R. Ma, Existence theorems for a second-order three-point boundary value problem, *J. Math. Anal. Appl.* **212** (1997), 430–442.
- [36] Yu. Melentsova, A best possible estimate of the nonoscillation interval for a linear differential equation with coefficients bounded in L_r , *Differ. Equ.* **13** (1977), 1236–1244.
- [37] Yu. Melentsova and G. Milshtein, An optimal estimate of the interval on which a multipoint boundary value problem possesses a solution, *Differ. Equ.* **10** (1974), 1257–1265.
- [38] Yu. Melentsova and G. Milshtein, Optimal estimation of the nonoscillation interval for linear differential equations with bounded coefficients, *Differ. Equ.* **17** (1981), 1368–1379.
- [39] S. K. Ntouyas and D. O'Regan, Existence results for semilinear neutral functional differential inclusions with nonlocal conditions, *Differ. Equ. Appl.* **1** (2009), no. 1, 41–65.
- [40] P. K. Palamides, G. Infante and P. Pietramala, Nontrivial solutions of a nonlinear heat flow problem via Sperner's lemma, *Appl. Math. Lett.* **22** (2009), no. 9, 1444–1450.
- [41] S. Roman and A. Štikonas, Greens functions for stationary problems with nonlocal boundary conditions, *Lith. Math. J.* **49** (2009), no. 2, 190–202.
- [42] H. B. Thompson and C. C. Tisdell, Three-point boundary value problems for second-order ordinary differential equations, *Math. Comput. Modelling* **34** (2001), 311–318.
- [43] J. Wang and Z. Zhang, Positive solutions to a second-order three-point boundary value problem, *J. Math. Anal. Appl.* **285** (2003), 237–249.
- [44] J. R. L. Webb, A unified approach to nonlocal boundary value problems, *Dynamic systems and applications* **5**, 510–515, Dynamic, Atlanta, GA, 2008.
- [45] J. R. L. Webb, Uniqueness of the principal eigenvalue in nonlocal boundary value problems, *Discrete Contin. Dyn. Syst. Ser. S* **1** (2008), no. 1, 177–186.
- [46] J. R. L. Webb, Remarks on nonlocal boundary value problems at resonance, *Appl. Math. Comput.* **216** (2010), no. 2, 497–500.
- [47] J. R. L. Webb and M. Zima, Multiple positive solutions of resonant and non-resonant nonlocal boundary value problems, *Nonlinear Anal.* **71** (2009), no. 3-4, 1369–1378.