


Title: Star Decompositions of the Complete Split Graph

Name: Adam Volk

Advisor: Atif Abueida, Ph.D.

Introduction


A graph is a discrete mathematical structure that consists of a set of vertices and a set of edges between pairs of vertices. A graph decomposition is a partitioning of the edges of a graph into disjoint sets in such a way that the induced subgraphs produced are isomorphic to each other. The graphs we focus on here are stars and complete split graphs (see below).



A complete split graph as the join of a complete graph and independent set

The Problem

Let G be the complete split graph with clique of order $n - m$ and independent set of order m . For what values of n , m , and t can we decompose G into edge disjoint copies of $K_{1,t}$?



Special Cases

- $m = n - 1$: decomposable if and only if $t \mid m$
- $t = 1$: trivial
- $t = 2$: decomposable if and only if total number of edges is even [3]

Necessary Conditions

- If G can be decomposed into t -stars, then $t \mid \binom{n-m}{2} + m(n-m)$

Casework and Results

- $n - m < t$: decomposable if and only if
 - $t \mid \frac{n+m-1}{2}$
- $n - m = t$: decomposable if and only if
 - t is odd and $m \geq \frac{t+1}{2}$
- $t < n - m < 2t$: decomposable if
 - $t \mid \frac{n+m-1}{2}$ or
 - t is odd, $t \mid m$, and $n - m = t + 1$
- $n - m \geq 2t$: decomposable if
 - $t \mid \frac{(n-m)(n-m-1)}{2}$ and $t \mid m(n-m)$,
 - $t \mid \frac{n+m-1}{2}$, or
 - $n - m$ is odd and $m \equiv -1 \pmod{t}$

Future work

Since we were unable to completely solve the problem for two of our cases, this is one place to begin.

We could also consider a more general problem by removing a subgraph H belonging to a different class of graphs.

Rather than limiting the size of stars to be a fixed value, we could consider decomposing a graph into stars of size t where t comes from some finite set of positive integers.

References

- [1] S.L. Hakimi, On the degrees of the vertices of a directed graph, J Franklin Inst 279 (1965), 290-308.
- [2] D.G. Hoffman and D. Roberts, Embedding partial k -star designs, Journal of Combinatorial Designs 22(4) (2014), 161-170
- [3] A. Kotzig, From the theory of finite regular graphs of degree three and four, Casopis Pest. Mat. 82 (1957), 7692
- [4] H.-C. Lee, Decomposition of the complete bipartite multigraph into cycles and stars, Discrete Mathematics 338 (2015), 1362-1369.
- [5] S. Yamamoto, H. Ikeda, S. Shige-eda, K. Ushio and N. Hamada, On claw-decomposition of complete graphs and complete bigraphs, Hiroshima Math. J. 5 (1975), 3342.